

Heisenberg and Modular Invariance of N=2 Conformal Field Theory

Shi-shyr Roan ¹

Institute of Mathematics

Academia Sinica

Taipei , Taiwan

(e-mail: maroan@ccvax.sinica.edu.tw)

Abstract

We present a theta function representation of the twisted characters for the rational N=2 superconformal field theory, and discuss the Jacobi-form like functional properties of these characters for a fixed central charge under the action of a finite Heisenberg group and modular transformations.

¹ Supported in part by the NSC grant of Taiwan.

1 Introduction

Superconformal algebras have been studied for a long time in physical literature. The importance is above all due to the fact that the superconformal algebra describes the underlying symmetries of Superstring theory. A superconformal algebra is a simple Lie superalgebra (over $\mathbb{C}$) spanned by modes of a finite family of local fields, which contains the Virasoro and some other odd, even ones, such that the coefficients of the operator product expansions are linear combinations of fields in the family and their derivatives. This infinite-dimensional algebra possesses a very rich mathematical structure with vast applications on different branches in physics and mathematics. Nevertheless, only until recent years, some rigorously systematic study of conformal superalgebras from the mathematical point of view has been worked out by groups of mathematicians and physicists, and profound mathematical structures have been found to further enhance our understanding of the original physical content (see, for instance [8, 13, 20] and references therein). However, despite the many theories of conformal algebras, the $N=2$ conformal theory has a peculiar nature which defines a class of its own, other than due to its applicable physical contents of the theory (see [4, 5, 6] references therein), also the mathematical reason on the intimate relation of special features of the $N=2$ superalgebra with Kähler geometry which has been a main core in the analytical study of projective algebraic manifolds. Hence it would be expected that one of the important aspects of $N=2$ conformal theory will be on the geometrical understanding of complex manifolds, especially the application on their topological invariants. A notable example would be the elliptic genus of $c_1 = 0$ manifolds, (see for instance [14, 21] and references therein), which amounts to the so called Gepner-model construction of the rational conformal theory [9]. As the mathematical structure of elliptic genus has now been well understood by works of the Hirzebruch school [11, 12], it is expected that a rigorous treatment of this topological invariant in terms of the $N=2$ conformal theory would be appeared as, or in a further complete form than, the one given in [18]. However, due to the length of presentation and also the nature of the contents, in this paper we shall only discuss the representation-theoretical aspect of the $N=2$ conformal algebra and leave its application on topology of manifolds as a separate problem, which will be treated in another work [19]. In this note, we shall study both the quantitative and qualitative properties of character functions of the rational $N=2$ conformal theory. The unitary $N=2$ superconformal characters have been analyzed by several groups of physicists [1, 3, 15, 16, 17]. While working on characters, they are usually regarded as the formal Laurent series; however a proper theta-form expression of the character is expected for the discussion of symmetries among the characters. By incorporating the characteristic of theta functions into the twisted currents of the algebra, we obtain a theta function (with characteristic) presentation of the characters of unitary irreducible highest weight modules (HWM) of $N=2$ conformal algebra for different sectors, (see Theorem 1 of the content). These explicit forms of characters in terms of the theta function, which seem to be unknown before to the best of the author's knowledge, do provide a convenient formulation for discussing the global functional properties through the classical elliptic function theory. Deceptively simple as it may appear, this is an crucial observation since it implies that in this form, the symmetries between characters of the rational $N=2$ conformal theory for a fixed central charge are clearly revealed and the later applications are expected to achieve in the study of the geometry of manifolds.

The paper is organized as follows. In Sect. 2, we give a brief introduction of the elliptic, modular functional properties of the Heisenberg and modular transformations. We also summarize a few known facts about the Jacobi-form like nature of theta functions (with characteristic) in the elliptic function theory. In Sect. 3, we first summarize some basic facts about the $N=2$ conformal algebra and the discrete series of unitary irreducible HWM with the central charge $c < 3$. Then we derive the theta form representation of characters of these HWM for all different

sectors, which enables us to further pursue their qualitative functional properties, instead of using only the Laurent series expression. In Sect. 4, we study the symmetries among these characters of HWM in the discrete series with a fixed central element. With the elliptic and modular properties, the characters constitute an irreducible representation of a central extension of some Jacobi-group, which composes of a finite Heisenberg group and a certain modular group. We consider this part as one of the most delicate aspects of this paper. For this reason we describe, as precisely as possible, the procedure of constructing the irreducible decomposition of Heisenberg group with a certain explicit basis, by which a matrix realization of modular transformations is given and the modular action on the irreducible Heisenberg-submodules is clearly shown. The detailed description will be stated in Theorem 2 of the paper. Finally, as it is known (cf., for instance [10] and references therein,) that superconformal algebras have a certain geometrical superspace realization in which generators are presented as super-vector fields (i.e., derivations) of super-spacetime. A such kind of geometrization for the usual Virasoro algebra has been a well-known process, by which the symmetry content of the algebra is better revealed. Hence in the last section as an appendix, we in particular recall and describe a detailed account of a geometrical realization of the N=2 conformal algebra as 1-dimensional vector fields with 2 supersymmetric extensions. This geometrical setting might offer a means of realizing the superconformal symmetry, which has been the focus of the present work.

Convention . In this paper, $\mathbb{R}, \mathbb{C}$ will denote the field of real, complex numbers respectively, and $\mathbb{H}$ the upper half plane $\{\tau \in \mathbb{C} \mid \text{Im}\tau > 0\}$; $\mathbb{R}^* = \mathbb{R} - \{0\}$, $\mathbb{C}^* = \mathbb{C} - \{0\}$. By a representation of a group G on a (complex) vector space V , we shall always mean a linear right-action of G on V ,

$$R : V \times G \longrightarrow V, \quad (v, g) \mapsto v \cdot g := R(v, g).$$

A R -eigenvector with the eigen-character ρ is an element $v \in V$ satisfying the relation:

$$v \cdot g = \rho(g)v, \quad \forall g \in G.$$

2 Elliptic and Modular Functional Properties

Denote $\mathcal{O}(\mathbb{C} \times \mathbb{H})$ the vector space of holomorphic functions of $\mathbb{C} \times \mathbb{H}$. In this section, we shall give a brief review on some basic elliptic and modular functional properties of $\mathbb{C} \times \mathbb{H}$, which will be needed for our later discussions. Here the ellipticity and modularity we refer to are the following standard actions of $\mathbb{R}^2, SL_2(\mathbb{R})$ on $\mathbb{C} \times \mathbb{H}$ respectively,

$$\begin{aligned} \mathbb{R}^2 \times (\mathbb{C} \times \mathbb{H}) &\longrightarrow \mathbb{C} \times \mathbb{H}, \quad (v, (z, \tau)) \mapsto (z + v_1\tau + v_2, \tau), \quad v = (v_1, v_2), \\ SL_2(\mathbb{R}) \times (\mathbb{C} \times \mathbb{H}) &\longrightarrow \mathbb{C} \times \mathbb{H}, \quad (M, (z, \tau)) \mapsto \left(\frac{z}{C\tau + D}, \frac{A\tau + B}{C\tau + D} \right), \quad M = \begin{pmatrix} A & B \\ C & D \end{pmatrix}. \end{aligned}$$

For the convenience of notations, in this section the letters, v_1, v_2 , will always denote the coordinates of a vector v in $\mathbb{R}^2$; A, B, C, D are the entries of a matrix M of $SL_2(\mathbb{R})$. Denote $\mathbb{R}^2 * SL_2(\mathbb{R})$ the semi-product of $\mathbb{R}^2$ and $SL_2(\mathbb{R})$ where the conjugation of $M \in SL_2(\mathbb{R})$ on $v \in \mathbb{R}^2$ is given by $M^{-1} \cdot v \cdot M = vM$ (the matrix product). The above expressions give rise to a geometrical action of $\mathbb{R}^2 * SL_2(\mathbb{R})$ on $\mathbb{C} \times \mathbb{H}$. One would like to lift this geometrical action to one on $\mathcal{O}(\mathbb{C} \times \mathbb{H})$. The elliptic part is as follows. Let $(\mathbb{R}^2, <, >)$ be the skew-symmetric form, $<v, v'> := v_1v'_2 - v_2v'_1$ for $v, v' \in \mathbb{R}^2$, and denote $\mathbb{C}_1^* = \{\alpha \in \mathbb{C} \mid |\alpha| = 1\}$. For $d \in \mathbb{R}^*$, $\mathbb{G}(d) = \mathbb{C}_1^* \times \mathbb{R}^2$ is the Heisenberg group (of index d) with the group law,

$$(\alpha, v) \cdot (\alpha', v') = (\alpha\alpha' e^{-2d\pi\sqrt{-1}v_2v'_1}, v + v'), \quad \text{for } \alpha, \alpha' \in \mathbb{C}_1^*, \quad v, v' \in \mathbb{R}^2.$$

We shall regard $\mathbb{C}_1^*, \mathbb{R}^2$ as the subsets of $\mathbb{G}(d)$ by the following identification of elements,

$$\mathbb{C}_1^* \ni \alpha := (\alpha, 0) \in \mathbb{G}(d) , \quad \mathbb{R}^2 \ni v := (1, v) \in \mathbb{G}(d) .$$

Then $\mathbb{G}(d)$ is generated by elements of $\mathbb{R}^2$ with the relations,

$$v \cdot v' = e^{2d\pi\sqrt{-1}\langle v, v' \rangle} v' \cdot v = e^{-2d\pi\sqrt{-1}v_2v'_1} (v + v') \in \mathbb{G}(d) , \quad \text{for } v, v' \in \mathbb{R}^2 .$$

As a character of $\mathbb{G}(d)$ is always trivial on its center, the character-group of $\mathbb{G}(d)$ is parametrized by the additive group $\mathbb{R}^2$, of which an element r corresponds the character ρ^r of $\mathbb{G}(d)$ with $\rho^r(v) := e^{2d\pi\sqrt{-1}\langle v, r \rangle}$ for $v \in \mathbb{R}^2 \subset \mathbb{G}(d)$. We shall denote $\Lambda(d)$ the subgroup of $\mathbb{G}(d)$ generated by integral elements in $\mathbb{R}^2$. When d is an integer, $\Lambda(d)$ is the abelian lattice group $\mathbb{Z}^2$, which will be denoted by Λ in what follows. Associated to an element τ of the upper half-plane $\mathbb{H}$, there is a $\mathbb{G}(d)$ -representation $\mathcal{T}(\tau; d)$ on the space of entire functions of $\mathbb{C}$,

$$\mathcal{T}(= \mathcal{T}(\tau; d)) : \mathcal{O}(\mathbb{C}) \times \mathbb{G}(d) \longrightarrow \mathcal{O}(\mathbb{C}) , \quad (\varphi, (\alpha, v)) \mapsto \varphi|_{\mathcal{T}_{(\alpha, v)}} := \alpha(\varphi|_{\mathcal{T}_v}) ,$$

where $(\varphi|_{\mathcal{T}_v})(z) := e^{d\pi\sqrt{-1}(v_1^2\tau + 2v_1(z+v_2))} \varphi(z + v_1\tau + v_2)$ for $v \in \mathbb{R}^2, z \in \mathbb{C}$. In particular, the following relations hold,

$$\mathcal{T}_{(v_1, v_2)} = \mathcal{T}_{(v_1, 0)} \mathcal{T}_{(0, v_2)} , \quad (\varphi|_{\mathcal{T}_{(v_1, 0)}})(z) = e^{d\pi\sqrt{-1}(v_1^2\tau + 2v_1z)} \varphi(z + v_1\tau) , \quad (\varphi|_{\mathcal{T}_{(0, v_2)}})(z) = \varphi(z + v_2) .$$

For $r \in \mathbb{R}^2$, the multiplication of the character ρ^r on $\mathcal{T}$ gives rise to a $\mathbb{G}(d)$ -representation $\mathcal{T}^r (= \mathcal{T}^r(\tau; d))$ on $\mathcal{O}(\mathbb{C})$,

$$(\varphi|_{\mathcal{T}_v^r})(z) = e^{2\pi\sqrt{-1}d\langle v, r \rangle} (\varphi|_{\mathcal{T}_v})(z) = e^{d\pi\sqrt{-1}(v_1^2\tau + 2v_1(z+v_2) + 2\langle v, r \rangle)} \varphi(z + v_1\tau + v_2) .$$

By $\mathcal{T}_r \mathcal{T}_*^r = \mathcal{T}_* \mathcal{T}_r$, $\mathcal{T}^r$ is equivalent to $\mathcal{T}$ via the intertwining operator $\mathcal{T}_r$. For $d \in \mathbb{R}^*$, $r \in \mathbb{R}^2$ and $\phi \in \mathcal{O}(\mathbb{C} \times \mathbb{H})$, we shall denote

$$\phi|_{\mathcal{T}^r} (= \phi|_{\mathcal{T}^r(d)}) := \phi(*, \tau)|_{\mathcal{T}^r(\tau; d)} \in \mathcal{O}(\mathbb{C} \times \mathbb{H}) . \quad (1)$$

The usual theta function theory is referred to the case $d = 1$. Recall the theta function $\vartheta(z, \tau)$ is an element in $\mathcal{O}(\mathbb{C} \times \mathbb{H})$ with the infinite product representation

$$\vartheta(z, \tau) = \prod_{n=1}^{\infty} (1 + 2q^{n-\frac{1}{2}} \cos 2\pi z + q^{2n-1})(1 - q^n) , \quad q = e^{2\pi\sqrt{-1}\tau} , \quad \text{Im}(\tau) \gg 0 ,$$

while the theta function with characteristics $r = (a, b) \in \mathbb{R}^2$ is defined by ²

$$\vartheta^r(z, \tau) := e^{\pi\sqrt{-1}(a^2\tau + 2a(z+b))} \vartheta(z + a\tau + b, \tau) , \quad (z, \tau) \in \mathbb{C} \times \mathbb{H} .$$

For a fixed τ , $\vartheta(z, \tau)$ (or $\vartheta^r(z, \tau)$) is the only Λ -invariant entire function on $\mathbb{C}$, up to a constant multiple, for the action $\mathcal{T}(1)$ ($\mathcal{T}^r(1)$ resp.) . In fact, one has the following expression of the theta functions,

$$\begin{aligned} \vartheta(z, \tau) &= (\mathbf{1} | \sum_{m=-\infty}^{\infty} \mathcal{T}_{(m, 0)}(1))(z, \tau) &= \sum_{m=-\infty}^{\infty} e^{\pi i(m^2\tau + 2mz)} , \\ \vartheta^r(z, \tau) &= ((\mathbf{1} | \mathcal{T}_r^r(1)) | \sum_{m=-\infty}^{\infty} \mathcal{T}_{(m, 0)}^r(1))(z, \tau) &= \sum_{m=-\infty}^{\infty} e^{\pi\sqrt{-1}((m+a)^2\tau + 2(m+a)(z+b))} , \end{aligned}$$

²Here we write the theta function $\vartheta^{(a, b)}$ instead of $\vartheta[\begin{smallmatrix} a \\ b \end{smallmatrix}]$ in usual literature.

where $\mathbf{1}$ is the constant function with value one. Associated to a theta function, one has the theta form,

$$\Theta(z, \tau) = \frac{\vartheta(z, \tau)}{\eta(\tau)^3}, \quad \Theta^r(z, \tau) = \frac{\vartheta^r(z, \tau)}{\eta(\tau)^3} \in \mathcal{O}(\mathbb{C} \times \mathbb{H}),$$

where $\eta(\tau) (= q^{1/24} \prod_{n=1}^{\infty} (1 - q^n))$ is the Dedekind eta function. The following relation holds,

$$\Theta^{(a,b)}(z, \tau) = e^{\pi\sqrt{-1}(a^2\tau+2a(z+\mu))} \Theta(z + a\tau + b, \tau), \quad (z, \tau) \in \mathbb{C} \times \mathbb{H}.$$

Corresponding to Jacobi theta function $\vartheta_1(z, \tau)$ ($:= \vartheta(\frac{1}{2}, \frac{1}{2})(z, \tau)$), we have the Jacobi-form

$$\Theta_1(z, \tau) := \Theta(\frac{1}{2}, \frac{1}{2})(z, \tau). \quad (2)$$

It is known that the asymptotic expansion of $\Theta_1(z, \tau)$ near $z = 0$ is given by

$$\Theta_1(z, \tau) = 2\pi z + O(z^2) \quad \text{as } z \rightarrow 0.$$

We have the quasi-periodicity and zero relations for $\Theta^{(a,b)}(z, \tau)$,

$$\begin{aligned} \Theta^{(a,b)}(z+1, \tau) &= e^{2\pi ia} \Theta^{(a,b)}(z, \tau), \quad \Theta^{(a,b)}(z+\tau, \tau) = e^{-\pi\sqrt{-1}(\tau+2(z+b))} \Theta^{(a,b)}(z, \tau), \\ \Theta^{(a,b)}(z, \tau) &= 0 \iff z \equiv (\tfrac{1}{2} - a)\tau + (\tfrac{1}{2} - b) \pmod{\mathbb{Z}\tau + \mathbb{Z}}. \end{aligned} \quad (3)$$

One has

$$\begin{aligned} \Theta^{(a+1,b)}(z, \tau) &= \Theta^{(a,b)}(z, \tau), \quad \Theta^{(a,b+1)}(z, \tau) = e^{2\pi\sqrt{-1}a} \Theta^{(a,b)}(z, \tau), \\ \Theta^{(a,b)}(-z, \tau) &= e^{2\pi\sqrt{-1}a} \Theta^{(1-a, 1-b)}(z, \tau), \\ \Theta^{(a+a', b+b')}(z, \tau) &= e^{\pi\sqrt{-1}(a^2\tau+2a(z+b+b'))} \Theta^{(a', b')}(z + a\tau + b, \tau). \end{aligned} \quad (4)$$

By lifting of the geometrical modular action, we have a $SL_2(\mathbb{R})$ -representation on $\mathcal{O}(\mathbb{C} \times \mathbb{H})$ for each pair (w, d) with $w \in \frac{1}{2}\mathbb{Z}, d \in \mathbb{R}^*$,

$$\begin{aligned} \Upsilon(= \Upsilon(w, d)) : \mathcal{O}(\mathbb{C} \times \mathbb{H}) \times SL_2(\mathbb{R}) &\longrightarrow \mathcal{O}(\mathbb{C} \times \mathbb{H}), \quad (\phi, M) \mapsto \phi|_{\Upsilon_M}, \\ (\phi|_{\Upsilon_M})(z, \tau) &:= (C\tau + D)^{-w} e^{\frac{-d\pi\sqrt{-1}Cz^2}{C\tau+D}} \phi\left(\frac{z}{C\tau+D}, \frac{A\tau+B}{C\tau+D}\right). \end{aligned} \quad (5)$$

The relation of the elliptic and modular actions on $\mathcal{O}(\mathbb{C} \times \mathbb{H})$ is given by the following lemma.

Lemma 1. For $(w, d) \in \frac{1}{2}\mathbb{Z} \times \mathbb{R}^*$, we denote $\mathcal{T}^r = \mathcal{T}^r(d), \Upsilon = \Upsilon(w, d)$. Then the following relation holds for $M \in SL_2(\mathbb{R}), v, v' \in \mathbb{R}^2$ with $v = v'M$, and for $\phi \in \mathcal{O}(\mathbb{C} \times \mathbb{H})$:

$$e^{-d\pi\sqrt{-1}(v_1v_2+2\langle v, r \rangle)} (\phi|_{\Upsilon_M})|_{\mathcal{T}_v^r} = e^{-d\pi\sqrt{-1}(v'_1v'_2+2\langle v', r \rangle)} (\phi|_{\mathcal{T}_{v'}^r})|_{\Upsilon_M}.$$

Proof. By the definition of $\mathcal{T}^r$, one needs only to consider the case $r = 0$. Note that $v_1 = v'_1A + v'_2C, v_2 = v'_1B + v'_2D$. By computation, one has the following relations:

$$\begin{aligned} (C\tau + D)^w ((\phi|_{\mathcal{T}_v^r})|_{\Upsilon_M})(z, \tau) &= e^{d\pi\sqrt{-1} \frac{-Cz^2+2v'_1z+(v_1'^2A+2v'_1v'_2C)\tau+v_1'^2B+2v'_1v'_2D}{C\tau+D}} \phi\left(\frac{z+v_1\tau+v_2}{C\tau+D}, \frac{A\tau+B}{C\tau+D}\right), \\ (C\tau + D)^w ((\phi|_{\Upsilon_M})|_{\mathcal{T}_v})(z, \tau) &= e^{d\pi\sqrt{-1} \left(\frac{-Cz^2+2v'_1z+(v_1'^2A+2v'_1v'_2C)\tau+v_1'^2B+2v'_1v'_2D}{C\tau+D} + v_1'^2AB+2v'_1v'_2BC+v_2'^2CD \right)} \\ &\quad \phi\left(\frac{z+v_1\tau+v_2}{C\tau+D}, \frac{A\tau+B}{C\tau+D}\right). \end{aligned}$$

By $v_1v_2 - v'_1v'_2 = v_1'^2AB + 2v'_1v'_2BC + v_2'^2CD$, the result follows immediately. $\square$

The conclusion of the above lemma has indicated that one can not lift the geometric action of $\mathbb{R}^2 * SL_2(\mathbb{R})$ of $\mathbb{C} \times \mathbb{H}$ directly to one of $\mathcal{O}(\mathbb{C} \times \mathbb{H})$. However, the formulas (1),(5) do give rise to a

module structure of $\mathcal{O}(\mathbb{C} \times \mathbb{H})$ for the semi-product $\mathbb{G}(d) \overset{r}{*} SL_2(\mathbb{R})$ of $\mathbb{G}(d)$ and $SL_2(\mathbb{R})$. Here in the group $\mathbb{G}(d) \overset{r}{*} SL_2(\mathbb{R})$, the conjugate relation of $M \in SL_2(\mathbb{R})$ on $v \in \mathbb{R}^2 \subseteq \mathbb{G}(d)$ is defined by

$$M \cdot v \cdot M^{-1} = e^{d\pi\sqrt{-1}(v_1v_2 - v'_1v'_2 + 2\langle v - v', r \rangle)} v' (= e^{d\pi\sqrt{-1}((v_1-2a)(v_2+2b) - (v'_1-2a)(v'_2+2b))} v') , \quad (6)$$

where $v' = (v'_1, v'_2)$ is the (matrix-product) element vM^{-1} . Note that $\mathbb{G}(d) \overset{r}{*} SL_2(\mathbb{R})$ is a central extension of $\mathbb{R}^2 \overset{r}{*} SL_2(\mathbb{R})$ with its center contained in $\mathbb{G}(d)$. Hence we have obtained the following result:

Proposition 1. The relations (1), (5) give rise to the representation of $\mathbb{G}(d) \overset{r}{*} SL_2(\mathbb{R})$ on $\mathcal{O}(\mathbb{C} \times \mathbb{H})$

$$\mathcal{J}^{d,r,w} : \mathcal{O}(\mathbb{C} \times \mathbb{H}) \times (\mathbb{G}(d) \overset{r}{*} SL_2(\mathbb{R})) \longrightarrow \mathcal{O}(\mathbb{C} \times \mathbb{H}). \quad (7)$$

□

For the rest of this section, we shall only consider the case $d \in \mathbb{Z}$, in which situation one will see that the representation (7) does descend to a certain arithmetic subgroup of $\mathbb{R}^2 \overset{r}{*} SL_2(\mathbb{R})$. Denote Γ the full modular group, $\Gamma = SL_2(\mathbb{Z})$. The subgroup $\Gamma^J := \Lambda \overset{r}{*} \Gamma$ of $\mathbb{R}^2 \overset{r}{*} SL_2(\mathbb{R})$ is called the Jacobi group. In general, the Jacobi group associated to a subgroup Γ' of Γ is defined by $\Gamma'^J := \Lambda' \overset{r}{*} \Gamma'$. It is easy to see that the following elements generates Γ^J ,

$$u = (1, 0), \quad v = (0, 1) \in \mathbb{Z}^2, \quad T = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, \quad S = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \in \Gamma.$$

In fact, Γ^J is characterized as the group generated by four elements u, v, T, S , with the relations,

$$\begin{aligned} uv &= vu, & S^4 &= 1, & S^2T &= TS^2, \\ T^{-1}uT &= u + v, & T^{-1}vT &= v, & S^{-1}uS &= v, & S^{-1}vS &= u^{-1}. \end{aligned}$$

One can easily see that a character λ of Γ^J is always trivial on the subgroup Λ , hence determined by the values of $\lambda(T), \lambda(S)$. Besides the full modular group Γ , the modular groups we shall also concern in this paper are the following one for $q = (q_1, q_2) \in \mathbb{Q}^2$,

$$\Gamma_q := \{M \in \Gamma \mid qM - q \in \Lambda\} (= \{M \in \Gamma \mid e^{2\pi\sqrt{-1}\langle vM - v, q \rangle} = 1 \ \forall v \in \Lambda\}) .$$

Note that Γ_q depends only on the class of q in $\mathbb{Q}^2/\Lambda$. The modular group $\Gamma_{(\frac{1}{2}, \frac{1}{2})}$ is generated by S, T^2 , which was denoted by Γ_θ in [18].

Proposition 2. Associated to an element r of $\mathbb{Q}^2$, we denote $q = (\frac{1}{2}, \frac{1}{2}) - r$. Then the following relation holds for $d, 2w \in \mathbb{Z}, v \in \Lambda$ and $M \in \Gamma_q$,

$$(\phi | \Upsilon_M(w, d)) | \mathcal{T}_{vM}^r(d) = (\phi | \mathcal{T}_v^r(d)) | \Upsilon_M(w, d), \quad \phi \in \mathcal{O}(\mathbb{C} \times \mathbb{H}).$$

Proof. Since the sign $(-1)^{(m+1)(n+1)}$ is unchanged under the action of Γ for $(m, n) \in \Lambda$, the above equality holds when $r = (\frac{1}{2}, \frac{1}{2})$. Then it follows the result for an arbitrary r by the definition of Γ_q . □

Remark. Applying the same argument in the proof of the above proposition to the representation (7) for $d \in \mathbb{Q}$, one obtains a $\Lambda(d) \overset{r}{*} \Gamma_q$ -module structure of $\mathcal{O}(\mathbb{C} \times \mathbb{H})$. □

For simplicity, we shall write $\phi|v, \phi|M$ instead of $\phi|\mathcal{T}_v^r(d), \phi|\Upsilon_M(w, d)$ in the case $d \in \mathbb{Z}$ if no confusion could arise. By Proposition 2, one obtains an Γ^J -representation on $\mathcal{O}(\mathbb{C} \times \mathbb{H})$, called the weight w and index $\frac{d}{2}$ representation of Γ^J ,

$$\begin{aligned} \mathcal{O}(\mathbb{C} \times \mathbb{H}) \times \Gamma_q^J &\longrightarrow \mathcal{O}(\mathbb{C} \times \mathbb{H}) , & (\phi, *) &\mapsto \phi|* , \\ (\phi|v)(z, \tau) &= e^{d\pi\sqrt{-1}(m^2\tau+2mz+2mb-2na)}\phi(z+m\tau+n, \tau) , & v &= (m, n) \in \Lambda , \\ (\phi|M)(z, \tau) &= (C\tau+D)^{-w}e^{\frac{-d\pi\sqrt{-1}Cz^2}{C\tau+D}}\phi\left(\frac{z}{C\tau+D}, \frac{A\tau+B}{C\tau+D}\right) , & M &\in \Gamma_q , \end{aligned} \quad (8)$$

where $q = (\frac{1}{2} - a, \frac{1}{2} - b)$. In particular, for $a = b = \frac{-1}{2}$ (or $\frac{1}{2}$), this is the usual weight w and index $\frac{d}{2}$ Γ^J -representation of which the invariant function $\phi(z, \tau)$ is called a (Γ^J) -Jacobi form of weight w and index $\frac{d}{2}$, i.e., the function $\phi(z, \tau)$ satisfies the following elliptic and modular properties [7]:

$$\begin{aligned} \phi(z+m\tau+n, \tau) &= e^{-d\pi\sqrt{-1}(m^2\tau+2mz-m+n)}\phi(z, \tau) , & (m, n) &\in \Lambda , \\ \phi\left(\frac{z}{C\tau+D}, \frac{A\tau+B}{C\tau+D}\right) &= (C\tau+D)^w e^{\frac{d\pi\sqrt{-1}Cz^2}{C\tau+D}}\phi(z, \tau) , & M &\in \Gamma . \end{aligned}$$

Denote $\mathcal{O}(\mathbb{C} \times \mathbb{H})_{w, \iota}$ the space of all Jacobi forms with weight w and index ι . Then with the canonical grading, the Jacobi forms form a graded algebras, $\bigoplus_{w, \iota \in \frac{1}{2}\mathbb{Z}} \mathcal{O}(\mathbb{C} \times \mathbb{H})_{w, \iota}$. An important example of Jacobi form is the Jacobi-theta form $\Theta_1(z, \tau)$ defined in (2), in fact, we have

$$\Theta_1(z, \tau) \in \mathcal{O}(\mathbb{C} \times \mathbb{H})_{-1, \frac{1}{2}} , \quad (9)$$

a conclusion followed by the facts that $\eta(\tau)$ is a weight $\frac{1}{2}$ Γ -modular eigenform for the eigencharacter σ , and $\vartheta_1(z, \tau)$ is a Jacobi eigenform of weight $\frac{1}{2}$ and index $\frac{1}{2}$ with the eigencharacter σ^3 , where σ is the character of Γ defined by $\sigma(T) = e^{\frac{\pi\sqrt{-1}}{12}}, \sigma(S) = e^{\frac{\pi\sqrt{-1}}{4}}$.

Remark. The discussion of this section on the Jacobi-form theory can be naturally extended to meromorphic forms on $\mathbb{C} \times \mathbb{H}$.

3 N=2 Superconformal Algebra

In this section, we shall derive the theta-form expression of characters of N=2 superconformal algebra representations. First we recall the some well-known facts on the algebra.

Definition. The N=2 superconformal algebra, denoted by SCA, consists of the stress tensor $L(\zeta)$, two super-currents $G^\pm(\zeta)$ and a $U(1)$ -current $J(\zeta)$,

$$L(\zeta) = \sum_{n \in \mathbb{Z}} L_n \zeta^{-n-2} , \quad G^\pm(\zeta) = \sum_{p \in \frac{1}{2} + \mathbb{Z}} G_p^\pm \zeta^{-p-\frac{3}{2}} , \quad J(\zeta) = \sum_{n \in \mathbb{Z}} J_n \zeta^{-n-1} ,$$

where the coefficients, $L_m, J_n, G_p^\pm$, form a super-Lie algebra with a central element c :

$$\begin{aligned} [L_m, L_n] &= (m-n)L_{m+n} + \frac{c(m^3-m)}{12}\delta_{m+n,0} , & [J_m, J_n] &= \frac{cm}{3}\delta_{m+n,0} , & [L_m, J_n] &= -nJ_{m+n} , \\ [L_m, G_p^\pm] &= \left(\frac{m}{2} - p\right)G_{m+p}^\pm , & [J_m, G_p^\pm] &= \pm G_{m+p}^\pm , \\ \{G_p^+, G_q^+\} &= \{G_p^-, G_q^-\} = 0 , & \{G_p^+, G_q^-\} &= 2L_{p+q} + (p-q)J_{p+q} + \frac{c}{3}(p^2 - \frac{1}{4})\delta_{p+q,0} . \end{aligned}$$

We shall not distinguish the central element c with its eigenvalues. For convenience, we shall use the bold letter $\mathbf{c}$ to denote one-third of the central element,

$$\mathbf{c} := \frac{c}{3} .$$

There exists the $\mathbb{Z}$ -symmetry of the algebra SCA, called the spectral flow, which is generated by the automorphism,

$$L_n \mapsto L_n + J_n + \frac{\mathbf{c}}{2}\delta_{n,0}, \quad J_n \mapsto J_n + \mathbf{c}\delta_{n,0}, \quad G_p^+ \mapsto G_{p+1}^+, \quad G_p^- \mapsto G_{p-1}^-.$$

In the topological field theory, it is more convenient to consider another set of generators of SCA associated to a real parameter $-\frac{1}{2} \leq a \leq \frac{1}{2}$ by the following twisting currents:

$$\begin{aligned} L^a(\zeta) &= L(\zeta) + a\partial J(\zeta) + \frac{\mathbf{c}a^2}{2}\zeta^{-2} = \sum_{n \in \mathbb{Z}} L_n^a \zeta^{-n-2}, \quad J^a(\zeta) = J(\zeta) + \mathbf{c}\zeta^{-1} = \sum_{n \in \mathbb{Z}} J_n^a \zeta^{-n-1}, \\ G^a(\zeta) &= G^+(\zeta) = \sum_{p \in -a+\frac{1}{2}+\mathbb{Z}} G_p^a \zeta^{-p-a-\frac{3}{2}}, \quad Q^a(\zeta) = G^-(\zeta) = \sum_{p \in a+\frac{1}{2}+\mathbb{Z}} Q_p^a \zeta^{-p+a-\frac{3}{2}}, \end{aligned}$$

with the coefficients

$$L_n^a = L_n + a(n+1)J_n + \frac{\mathbf{c}a^2}{2}\delta_{n,0}, \quad J_n^a = J_n + \mathbf{c}a\delta_{n,0}, \quad G_p^a = G_{p+a}^+, \quad Q_p^a = G_{p-a}^-,$$

satisfying the relations,

$$\begin{aligned} [L_m^a, L_n^a] &= (m-n)L_{m+n}^a + \frac{\mathbf{c}(m^3(1-4a^2)-m)}{4}\delta_{m+n,0}, & [J_m^a, J_n^a] &= \mathbf{c}m\delta_{m+n,0}, \\ [L_m^a, J_n^a] &= -nJ_{m+n}^a + \mathbf{c}n^2\delta_{m+n,0}, & [L_m^a, Q_p^a] &= (\frac{(1-2a)m}{2} - p)Q_{m+p}^a, \\ [L_m^a, G_p^a] &= (\frac{(1+2a)m}{2} - p)G_{m+p}^a, & \{G_p^a, G_q^a\} &= \{Q_p^a, Q_q^a\} = 0, \\ [J_m^a, G_p^a] &= G_{m+p}^a, \quad [J_m^a, Q_p^a] = -Q_{m+p}^a, & \{G_p^a, Q_q^a\} &= \{Q_p^a, Q_q^a\} = 0, \\ \{G_p^a, Q_q^a\} &= 2L_{p+q}^a + ((1-2a)p - (1+2a)q)J_{p+q}^a + \frac{\mathbf{c}(4p^2-1)}{4}\delta_{p+q,0}. \end{aligned}$$

One has the decomposition, $\text{SCA} = \text{SCA}_+ \oplus \text{SCA}_0 \oplus \text{SCA}_-$, where

$$\begin{aligned} \text{SCA}_0 &= \langle c, L_0, J_0 \rangle = \langle c, L_0^a, J_0^a \rangle, \\ \text{SCA}_+ &= \langle L_m, J_m, G_p^\pm \mid m > 0, p \geq 0 \rangle = \langle L_m^a, J_m^a, G_p^a, Q_p^a \mid m > 0, p \geq 0 \rangle, \\ \text{SCA}_- &= \langle L_m, J_m, G_p^\pm \mid m < 0, p \leq 0 \rangle = \langle L_m^a, J_m^a, G_p^a, Q_p^a \mid m < 0, p \leq 0 \rangle. \end{aligned}$$

A highest weight module (HWM) is a SCA-module generated by a highest vector, i.e., an eigenvector of SCA_0 and annihilated by SCA_+ . The letters H, Q will denote the L_0, J_0 -eigenvalues of the highest vector respectively; similarly the letters H^a, Q^a for L_0^a, J_0^a -eigenvalues. Then $H^a = H + aQ + \frac{a^2\mathbf{c}}{2}$, $Q^a = Q + a\mathbf{c}$. The character of a HWM is the following Laurent series,

$$\text{NS}(z, \tau) = \text{Tr}(q^{L_0 - \frac{\mathbf{c}}{8}} y^{J_0}) \quad , \quad y := e^{2\pi\sqrt{-1}z}, q := e^{2\pi\sqrt{-1}\tau}.$$

The twisted character with characteristic (a, b) for $-\frac{1}{2} \leq a, b \leq \frac{1}{2}$ is defined by

$$\text{Ch}^{(a,b)}(z, \tau) = \text{Tr}(q^{L_0 - \frac{\mathbf{c}}{8}} (e^{2\pi\sqrt{-1}bz})^{J_0^a}) = e^{\mathbf{c}\pi\sqrt{-1}(a^2\tau + 2a(z+b))} \text{NS}(z + a\tau + b, \tau) \quad (10)$$

By

$$\begin{aligned} L_0^{1-a} &= L_0^a + (1-2a)J_0^a + \frac{(1-2a)^2\mathbf{c}}{2}, & J_0^{1-a} &= J_0^a + (1-2a)\mathbf{c}; \\ L_0^{\frac{1}{2}} &= L_0^{-\frac{1}{2}} + J_0^{-\frac{1}{2}} + \frac{\mathbf{c}}{2}, & J_0^{\frac{1}{2}} &= J_0^{-\frac{1}{2}} + \mathbf{c}, \end{aligned}$$

one obtains the following relations of twisted characters,

$$\begin{aligned} \text{Ch}^{(1-a,b)}(z, \tau) &= e^{\mathbf{c}\pi\sqrt{-1}((1-2a)^2\tau + 2(1-2a)(z+b))} \text{Ch}^{(a,b)}(z + (1-2a)\tau, \tau), \\ \text{Ch}^{(a,1-b)}(z + (1-2b)\tau, \tau) &= \text{Ch}^{(a,b)}(z, \tau), \\ \text{Ch}^{(\frac{1}{2},b)}(z, \tau) &= e^{\mathbf{c}\pi\sqrt{-1}(\tau + 2(z+b))} \text{Ch}^{(\frac{-1}{2},b)}(z + \tau, \tau), \quad \text{Ch}^{(a,\frac{1}{2})}(z, \tau) = \text{Ch}^{(a,\frac{-1}{2})}(z + 1, \tau). \end{aligned} \quad (11)$$

We shall denote³

$$R(z, \tau) = \text{Ch}^{(\frac{1}{2}, \frac{1}{2})}(z, \tau) , \quad \tilde{R}(z, \tau) = \text{Ch}^{(\frac{-1}{2}, \frac{-1}{2})}(z, \tau) .$$

For $0 < \mathbf{c} < 1$, all the unitary irreducible HWM of SCA are labelled by three integers k, m, l with the relations, $|m| \leq l \leq k$, $l \equiv m \pmod{2}$, where $\mathbf{c}, H, Q$ are given by

$$\mathbf{c} = \frac{k}{k+2} , \quad H(= H_{l,m}) = \frac{l^2+2l-m^2}{4(k+2)} , \quad Q(= Q_{l,m}) = \frac{m}{k+2} . \quad (12)$$

The character of the corresponding HWM will be denoted by $\text{Ch}_{l,m}^{(a,b)}(z, \tau)$. A HWM is called chiral (resp. antichiral) if $H_{l,m} = \frac{Q_{l,m}}{2}$ (resp. $\frac{-Q_{l,m}}{2}$), characterized by the highest vector annihilated by $G_{\frac{-1}{2}}^+$ (resp. $G_{\frac{-1}{2}}^-$) . For the rest of this paper, associated to a positive integer k we shall use the bold letter $\mathbf{k}$ to denote the number $k+2$,

$$\mathbf{k} := k + 2 .$$

To a pair of integers l, m with $l \equiv m \pmod{2}$, we shall always use the letters i, j to denote the following half-integers :

$$i = \frac{l-m+1}{2} , \quad j = \frac{-(l+m+1)}{2} \in \frac{1}{2} + \mathbb{Z} .$$

With a fixed central element $\mathbf{c} = \frac{\mathbf{k}-2}{\mathbf{k}}$, the unitary irreducible HWMs are indexed by the following equivalent data,

$$(l, m) \in \mathbb{Z}^2 , \quad |m| \leq l \leq k , \quad l \equiv m \pmod{2} \iff (i, j) \in (\frac{1}{2} + \mathbb{Z})^2 , \quad 0 < i, -j, i-j < \mathbf{k} ,$$

hence the chirality is given by

$$\begin{aligned} \text{chiral HWM :} \quad & m = l, \quad \iff \quad i = \frac{1}{2} , \\ \text{antichiral HWM :} \quad & m = -l, \quad \iff \quad j = \frac{-1}{2} . \end{aligned}$$

Then (12) becomes

$$\begin{aligned} H &= \frac{-4ij-1}{4\mathbf{k}} , \quad H^a = \frac{l^2+2l-(m-2a)^2}{4\mathbf{k}} + \frac{a^2}{2} = \frac{-4(i+a)(j+a)-1}{4\mathbf{k}} + \frac{a^2}{2} , \\ Q &= \frac{-(i+j)}{\mathbf{k}} , \quad Q^a = \frac{m+a(\mathbf{k}-2)}{\mathbf{k}} = \frac{-(i+j)+a(\mathbf{k}-2)}{\mathbf{k}} . \end{aligned} \quad (13)$$

For the rest of this section, we are going to derive the theta-form expression of the character of the above HWM and discuss the functional properties among these characters. First let us recall the Laurent series expression⁴ of $\text{NS}_{l,m}(z, \tau)$,

$$\text{NS}_{l,m}(z, \tau) = q^{H-\frac{\mathbf{c}}{8}} y^Q \gamma_{l,m}(z, \tau) \varphi(z, \tau) ,$$

where

$$\begin{aligned} \varphi(z, \tau) &= \frac{\prod_{n=1}^{\infty} (1+yq^{n-1/2})(1+y^{-1}q^{n-1/2})}{\prod_{n=1}^{\infty} (1-q^n)^2} , \\ \gamma_{l,m}(z, \tau) &= \frac{\prod_{n=1}^{\infty} (1-q^{\mathbf{k}(n-1)+i-j})(1-q^{\mathbf{k}n-i+j})(1-q^{\mathbf{k}n})^2}{\prod_{n=1}^{\infty} (1+yq^{\mathbf{k}(n-1)+i})(1+y^{-1}q^{\mathbf{k}n-i})(1+yq^{\mathbf{k}n+j})(1+y^{-1}q^{\mathbf{k}(n-1)-j})} . \end{aligned}$$

³The letters, NS, R, indicate the Neveu-Schwarz or Ramond sectors.

⁴ Here we adhere to papers [3, 16, 17] for the mostly suitable format to our treatment.

Theorem 1. The characters $\text{NS}_{l,m}(z, \tau)$, $\text{Ch}_{l,m}^{(a,b)}(z, \tau)$ are elements in $\mathcal{O}(\mathbb{C} \times \mathbb{H})$ with the following theta-form representation:

$$\begin{aligned} \text{NS}_{l,m}(z, \tau) &= \frac{e^{\pi\sqrt{-1}(\frac{1}{2}-\frac{i-j}{k})}\Theta(\frac{1}{2}+\frac{i-j}{k}, \frac{1}{2})(0, \mathbf{k}\tau)\Theta(z, \tau)}{\Theta(\frac{1}{2}+\frac{i}{k}, 0)(z, \mathbf{k}\tau)\Theta(\frac{1}{2}+\frac{j}{k}, 0)(z, \mathbf{k}\tau)}, \\ \text{Ch}_{l,m}^{(a,b)}(z, \tau) &= \frac{e^{\pi\sqrt{-1}(\frac{1}{2}-\frac{i-j}{k})}\Theta(\frac{1}{2}+\frac{i-j}{k}, \frac{1}{2})(0, \mathbf{k}\tau)\Theta^{(a,b)}(z, \tau)}{\Theta(\frac{1}{2}+\frac{i+a}{k}, b)(z, \mathbf{k}\tau)\Theta(\frac{1}{2}+\frac{j+a}{k}, b)(z, \mathbf{k}\tau)}. \end{aligned} \quad (14)$$

Proof. By the infinite product formula of $\vartheta(z, \tau)$, the factors $\varphi(z, \tau)$, $\gamma_{l,m}(z, \tau)$ in the formula of $\text{NS}(z, \tau)$ have the following expressions:

$$\varphi(z, \tau) = q^{\frac{1}{8}}\Theta(z, \tau), \quad \gamma_{l,m}(z, \tau) = \frac{q^{-\frac{k}{8}}\Theta((\frac{1}{2}-\frac{i-j}{k})\mathbf{k}\tau + \frac{1}{2}, \mathbf{k}\tau)}{\Theta(z + (\frac{-1}{2} + \frac{i}{k})\mathbf{k}\tau, \mathbf{k}\tau)\Theta(z + (\frac{1}{2} + \frac{j}{k})\mathbf{k}\tau, \mathbf{k}\tau)}.$$

By (13) (4), we have

$$\gamma_{l,m}(z, \tau) = \frac{q^{-H+\frac{c-1}{8}}y^{-Q}e^{\pi\sqrt{-1}(\frac{-1}{2}+\frac{i-j}{k})}\Theta(\frac{1}{2}-\frac{i-j}{k}, \frac{1}{2})(0, \mathbf{k}\tau)}{\Theta(\frac{-1}{2}+\frac{i}{k}, 0)(z, \mathbf{k}\tau)\Theta(\frac{1}{2}+\frac{j}{k}, 0)(z, \mathbf{k}\tau)} = \frac{q^{-H+\frac{c-1}{8}}y^{-Q}e^{\pi\sqrt{-1}(\frac{1}{2}-\frac{i-j}{k})}\Theta(\frac{1}{2}+\frac{i-j}{k}, \frac{1}{2})(0, \mathbf{k}\tau)}{\Theta(\frac{1}{2}+\frac{i}{k}, 0)(z, \mathbf{k}\tau)\Theta(\frac{1}{2}+\frac{j}{k}, 0)(z, \mathbf{k}\tau)}.$$

Hence we obtain the expression of $\text{NS}_{l,m}(z, \tau)$. By the last relation of (4), it follows the expression of $\text{Ch}_{l,m}^{(a,b)}(z, \tau)$. The zeros of the denominator in the expression of $\text{Ch}_{l,m}^{(a,b)}(z, \tau)$ are given by

$$z \equiv -(i+a)\tau + \frac{1}{2} - b, \quad -(j+a)\tau + \frac{1}{2} - b \pmod{\mathbb{Z}\mathbf{k}\tau + \mathbb{Z}}.$$

By $0 < i - j < k$, there is no common solution for the above two equations. Hence the poles of $\text{Ch}_{l,m}^{(a,b)}(z, \tau)$ are all simple; so are the zeros of $\Theta^{(a,b)}(z, \tau)$ in the numerator of the expression. Then one can easily see that $\text{Ch}_{l,m}^{(a,b)}(z, \tau) \in \mathcal{O}(\mathbb{C} \times \mathbb{H})$. $\square$

By the last relation of (4) for $(a, b) = \pm(\frac{1}{2}, \frac{1}{2})$, the Ramon characters are expressed by

$$\begin{aligned} \text{R}_{l,m}(z, \tau) &= \frac{\sqrt{-1}q^{H+\frac{Q}{2}}(-y)^{Q+\frac{c-1}{2}}\Theta_1((i-j)\tau, \mathbf{k}\tau)\Theta_1(z, \tau)}{\Theta_1(z+(i+\frac{1}{2})\tau, \mathbf{k}\tau)\Theta_1(z+(j+\frac{1}{2})\tau, \mathbf{k}\tau)}, \\ \tilde{\text{R}}_{l,m}(z, \tau) &= \frac{e^{\frac{-\pi\sqrt{-1}}{2}}q^{H-\frac{Q}{2}}(e^{-\pi\sqrt{-1}}y)^{Q-\frac{c-1}{2}}\Theta_1((i-j)\tau, \mathbf{k}\tau)\Theta_1(z, \tau)}{\Theta_1(z+(i-\frac{1}{2})\tau, \mathbf{k}\tau)\Theta_1(z+(j-\frac{1}{2})\tau, \mathbf{k}\tau)}. \end{aligned} \quad (15)$$

By (4) (11), and the relations

$$1 \leq i + \frac{1}{2} \leq k + 1, \quad -k \leq j + \frac{1}{2} \leq 0, \quad 0 \leq i - \frac{1}{2} \leq k, \quad, \quad -(k+1) \leq j - \frac{1}{2} \leq -1,$$

one obtains the following relations of Ramon characters, which was used in the study of topological elliptic genus [6, 18]

Corollary 1.⁵

$$\begin{aligned}
R_{l,-m}(z, \tau) &= \tilde{R}_{l,m}(-z, \tau), & \tilde{R}_{l,-m}(z, \tau) &= R_{l,m}(-z, \tau), \\
R_{l,m}(-z, \tau) &= e^{c\pi\sqrt{-1}(\tau-2z+1)} R_{l,-m}(z - \tau - 1, \tau), & \tilde{R}_{l,m}(-z, \tau) &= e^{c\pi\sqrt{-1}(\tau+2z+1)} \tilde{R}_{l,-m}(z + \tau + 1, \tau), \\
R_{l,-m}(z, \tau) &= e^{c\pi\sqrt{-1}(\tau+2z+1)} R_{l,m}(-z - \tau - 1, \tau), & \tilde{R}_{l,-m}(z, \tau) &= e^{c\pi\sqrt{-1}(\tau-2z+1)} \tilde{R}_{l,m}(-z + \tau + 1, \tau), \\
R_{l,m}(0, \tau) &= \begin{cases} (-1)^{Q_{l,m}+\frac{c}{2}} & \text{if } m = -l, \text{ i.e. antichiral,} \\ 0 & \text{otherwise,} \end{cases} \\
\tilde{R}_{l,m}(0, \tau) &= \begin{cases} (-1)^{-Q_{l,m}+\frac{c}{2}} & \text{if } m = l, \text{ i.e. chiral,} \\ 0 & \text{otherwise.} \end{cases}
\end{aligned}$$

□

For characters of other sectors, by (3) (4) and Theorem 1, one has the following result.

Corollary 2. The following relations hold for characters of HWMs for a fixed central element $\mathbf{c}$:

$$e^{2\pi\sqrt{-1}Q_{l,m}} \text{Ch}_{l,m}^{(a,b)}(z, \tau) = e^{-2c\pi\sqrt{-1}a} \text{Ch}_{l,m}^{(a,b)}(z+1, \tau), \quad \text{Ch}_{l,m-2}^{(a,b)}(z, \tau) = e^{c\pi\sqrt{-1}(\tau+2(z+b))} \text{Ch}_{l,m}^{(a,b)}(z+\tau, \tau).$$

□

For the qualitative study of characters, it is more convenient to extend the domain of (l, m) to the following larger one:

$$\mathcal{L}_{\mathbf{c}} := \{(l, m) \in \mathbb{Z}^2 \mid 0 \leq l \leq \mathbf{k} - 2, l \equiv m \pmod{2}\} \left(\Longleftrightarrow \{(i, j) \in (\frac{1}{2} + \mathbb{Z})^2 \mid 0 < i - j < \mathbf{k}\} \right).$$

For $(l, m) \in \mathcal{L}_{\mathbf{c}}$, one defines $H_{l,m}, Q_{l,m}, \text{Ch}_{l,m}^{(a,b)}(z, \tau)$ etc. by the same expressions in (12) (13), (14), (15). The chirality now becomes,

$$\text{chiral} \Longleftrightarrow e^{2\pi\sqrt{-1}(H_{(l,m)} - \frac{Q_{(l,m)}}{2})} = 1, \quad \text{antichiral} \Longleftrightarrow e^{2\pi\sqrt{-1}(H_{(l,m)} + \frac{Q_{(l,m)}}{2})} = 1$$

Define the automorphism of $\mathcal{L}_{\mathbf{c}}$,


$$r : \mathcal{L}_{\mathbf{c}} \longrightarrow \mathcal{L}_{\mathbf{c}}, \quad (l, m) \mapsto (\mathbf{k} - 2 - l, m + \mathbf{k}) \left(\Longleftrightarrow (i, j) \mapsto (j, i) - (0, \mathbf{k}) \right).$$

Denote $\mathcal{B}_{\mathbf{c}}$ be the space of $< r >$ -orbits, $\mathcal{V}_{\mathbf{c}}$ the Hilbert space with $\mathcal{B}_{\mathbf{c}}$ as an orthonormal basis,

$$\mathcal{B}_{\mathbf{c}} = \mathcal{L}_{\mathbf{c}} / < r >, \quad \mathcal{V}_{\mathbf{c}} = \oplus_{\lambda \in \mathcal{B}_{\mathbf{c}}} \mathbb{C}\lambda.$$

A fundamental region for $\mathcal{B}_{\mathbf{c}}$ consists of those (l, m) with $|m| \leq l$, i.e., the index set of HWM.

⁵ In the paper [18], the signs in the formula of $\tilde{R}_{\lambda}$ on pp. 373 were inaccurate in order to keep the statement of Proposition 4 there to be valid. The correct one should be $\tilde{R}_{\lambda}(z, \tau) = NS_{\lambda}(z - \frac{\tau}{2} - \frac{1}{2}, \tau) \exp(\frac{c(K)}{3}\pi\sqrt{-1}(\frac{\tau}{4} - z + 1/2))$. Then Lemma 4 in [18] holds for R_{λ} of this present paper, instead of $\tilde{R}_{\lambda}$.



Note that the characters and chirality thus defined are invariant under the automorphism r , i.e., depending only on the class of (l, m) in $\mathcal{B}_c$, which will be denoted by $[l, m]$. We denote

$$\text{Ch}_\lambda^{(a,b)}(z, \tau) := \text{Ch}_{l,m}^{(a,b)}(z, \tau) \quad \text{for } \lambda = [l, m] \in \mathcal{B}_c,$$

similar for $\text{NS}_\lambda(z, \tau)$, $\text{R}_\lambda(z, \tau)$, $\tilde{\text{R}}_\lambda(z, \tau)$. We introduce the following automorphisms of $\mathcal{L}_c$,

$$\begin{aligned} f_0 : (l, m) &\mapsto (l, m - 2) & (\iff (i, j) &\mapsto (i + 1, j + 1)) , \\ n_0 : (l, m) &\mapsto (l, -m) & (\iff (i, j) &\mapsto (-j, -i)) . \end{aligned}$$

By $f_0 r = r f_0$, $n_0 r = r^{-1} n_0$, the maps f_0, n_0 induce the one-one correspondences of $\mathcal{B}_c$, hence the linear automorphisms f, n of the vector space $\mathcal{V}_c$ with the order $k, 2$ respectively,

$$\begin{aligned} f : \mathcal{V}_c &\longrightarrow \mathcal{V}_c, & [l, m] \in \mathcal{B}_c &\mapsto [l, m - 2] \in \mathcal{B}_c, \\ n : \mathcal{V}_c &\longrightarrow \mathcal{V}_c, & [l, m] \in \mathcal{B}_c &\mapsto [l, -m] \in \mathcal{B}_c. \end{aligned} \tag{16}$$

The linear transformations f, n are called the spectral flow, the charge conjugation of $\mathcal{V}_c$ respectively. By Corollary 1 of Theorem 1, one has the following relations of Ramon characters:

$$\begin{aligned} \text{R}_{n(\lambda)}(z, \tau) &= \tilde{\text{R}}_\lambda(-z, \tau), & \tilde{\text{R}}_{n(\lambda)}(z, \tau) &= \text{R}_\lambda(-z, \tau), \\ \text{R}_\lambda(-z, \tau) &= e^{c\pi\sqrt{-1}(\tau-2z+1)} \text{R}_{n(\lambda)}(z - \tau - 1, \tau), & \tilde{\text{R}}_\lambda(-z, \tau) &= e^{c\pi\sqrt{-1}(\tau+2z+1)} \tilde{\text{R}}_{n(\lambda)}(z + \tau + 1, \tau), \\ \text{R}_{n(\lambda)}(z, \tau) &= e^{c\pi\sqrt{-1}(\tau+2z+1)} \text{R}_\lambda(-z - \tau - 1, \tau), & \tilde{\text{R}}_{n(\lambda)}(z, \tau) &= e^{c\pi\sqrt{-1}(\tau-2z+1)} \tilde{\text{R}}_\lambda(-z + \tau + 1, \tau). \end{aligned} \tag{17}$$

By Corollary 2 of Theorem 1, we have the following relations of $\text{Ch}_\lambda^{(a,b)}(z, \tau)$ for $\lambda \in \mathcal{B}_c$, by which the elliptic property of characters will be discussed in the next section,

$$e^{c\pi\sqrt{-1}(\tau+2(z+b))} \text{Ch}_\lambda^{(a,b)}(z + \tau, \tau) = \text{Ch}_{f(\lambda)}^{(a,b)}(z, \tau), \quad e^{-2c\pi\sqrt{-1}a} \text{Ch}_\lambda^{(a,b)}(z + 1, \tau) = e^{2\pi\sqrt{-1}Q_\lambda} \text{Ch}_\lambda^{(a,b)}(z, \tau), \tag{18}$$

in particular, for the Neveu-Schwarz and Ramon characters one has

$$\begin{aligned} e^{c\pi\sqrt{-1}(\tau+2z)}\text{NS}_\lambda(z+\tau,\tau) &= \text{NS}_{f(\lambda)}(z,\tau) , & \text{NS}_\lambda(z+1,\tau) &= e^{2\pi\sqrt{-1}Q_\lambda}\text{NS}_\lambda(z,\tau) , \\ e^{c\pi\sqrt{-1}(\tau+2z+1)}\text{R}_\lambda(z+\tau,\tau) &= \text{R}_{f(\lambda)}(z,\tau), & e^{-c\pi\sqrt{-1}}\text{R}_\lambda(z+1,\tau) &= e^{2\pi\sqrt{-1}Q_\lambda}\text{R}_\lambda(z,\tau) , \\ e^{c\pi\sqrt{-1}(\tau+2z-1)}\tilde{\text{R}}_\lambda(z+\tau,\tau) &= \tilde{\text{R}}_{f(\lambda)}(z,\tau) , & e^{c\pi\sqrt{-1}}\tilde{\text{R}}_\lambda(z+1,\tau) &= e^{2\pi\sqrt{-1}Q_\lambda}\tilde{\text{R}}_\lambda(z,\tau) . \end{aligned} \quad (19)$$

4 Ellipticity and Modularity of Superconformal Algebra Representations

In this section, we are going to discuss the symmetry properties among the unitary irreducible HWMs with a fixed central element $\mathbf{c} = \frac{\mathbf{k}-2}{\mathbf{k}}$ for SCA-modules. First we describe some notions which will facilitate the later presentation significantly. For a positive integer N , a vector v in $\mathfrak{C}^N$ will be identified with a sequence of complex numbers, $(v_m)_{m \in \mathbb{Z}}$, with $v_m = v_{m+N}$. The standard basis of $\mathfrak{C}^N$ are denoted by $|n\rangle, n \in \mathbb{Z}$ with $|n\rangle = |m\rangle$ if $n \equiv m \pmod{N}$. Denote V_N, U_N and U_N^ζ ($\zeta \in \mathfrak{C}^*$) as the linear automorphisms of $\mathfrak{C}^N$ defined by

$$\begin{aligned} V_N|n\rangle &= \omega^n|n\rangle , & U_N|n\rangle &= |n+1\rangle , \\ U_N^\zeta|n\rangle &= \begin{cases} \zeta|n+1\rangle & \text{if } n \equiv 0 \pmod{N}, \\ |n+1\rangle & \text{otherwise.} \end{cases} \end{aligned}$$

Here ω is a primitive N -th root of unity, which we will specify later.

For a given $r = (a, b)$, with the identification of the basis element λ of $\mathcal{V}_{\mathbf{c}}$ and the character $\text{Ch}_\lambda^{(a,b)}(z, \tau)$, the relation (18) gives rise to a $\Lambda(\mathbf{c})$ -representation on $\mathcal{V}_{\mathbf{c}}$ which describes the elliptic properties of the HWMs. This $\Lambda(\mathbf{c})$ -representation is trivial on the subgroup $\mathbf{k}\Lambda$ of $\Lambda(\mathbf{c})$ which is generated by $(\mathbf{k}, 0)$ and $(0, \mathbf{k})$, hence it descends to a representation of the finite quotient group, $\overline{\Lambda(\mathbf{c})} := \Lambda(\mathbf{c})/\mathbf{k}\Lambda$,

$$\mathcal{V}_{\mathbf{c}} \times \overline{\Lambda(\mathbf{c})} \longrightarrow \mathcal{V}_{\mathbf{c}} . \quad (20)$$

For easier description of the above representation, we define a pair of integers N, M associated to the central element $\mathbf{c} = \frac{\mathbf{k}-2}{\mathbf{k}}$,

$$(N, M) = (N(\mathbf{c}), M(\mathbf{c})) = \begin{cases} (\mathbf{k}, \frac{\mathbf{k}-1}{2}) & \text{for odd } \mathbf{k}, \\ (\frac{\mathbf{k}}{2}, \mathbf{k}-1) & \text{for even } \mathbf{k}. \end{cases}$$

Denote $P = \overline{(1, 0)}, Q = \overline{(0, 1)}$, the generators of $\overline{\Lambda(\mathbf{c})}$, and they satisfy the Weyl commutation relation:

$$PQ = \omega^{-1}QP , \quad \omega := e^{-2\pi\sqrt{-1}\mathbf{c}} = e^{\frac{4\pi\sqrt{-1}}{\mathbf{k}}} \quad (21)$$

Note that ω is a primitive N -th root of unity. It is known that the irreducible unitary representations of $\overline{\Lambda(\mathbf{c})}$ are parametrized by elements (ζ_1, ζ_2) in the (real) 2-torus $\mathfrak{C}_1^* \times \mathfrak{C}_1^*$, where (ζ_1, ζ_2) represents the N -dimensional $\Lambda(\mathbf{c})$ -module $\mathfrak{C}_{(\zeta_1, \zeta_2)}^N$ with $Q \mapsto \zeta_1^{\frac{1}{N}}V_N, P \mapsto \zeta_2^{\frac{1}{N}}U_N$. Note that $\mathfrak{C}_{(\zeta_1, \zeta_2)}^N$ is equivalent to the representation with $Q \mapsto \zeta_1^{\frac{1}{N}}V_N, P \mapsto U_N^{\zeta_2}$.

Lemma 2. The $\overline{\Lambda(\mathbf{c})}$ -module $\mathcal{V}_{\mathbf{c}}$ is the following direct sum of M irreducible representations,

$$\mathcal{V}_{\mathbf{c}} = \begin{cases} M\mathfrak{C}_{(1,1)}^N & \text{for odd } \mathbf{k}, \\ \bigoplus_{\epsilon, \epsilon' = \pm 1} M_{\epsilon, \epsilon'}\mathfrak{C}_{(\epsilon, \epsilon')}^N & \text{for even } \mathbf{k}, \end{cases}$$

where the multiplicities $M_{\epsilon, \epsilon'}$ for even $\mathbf{k}$ are defined by

$$\begin{cases} M_{\epsilon, \epsilon'} = \frac{\mathbf{k}-2}{4} & \text{except } M_{1,1} = \frac{\mathbf{k}+2}{4}, \quad \text{for } \frac{\mathbf{k}}{2} \equiv 1 \pmod{2}, \\ M_{\epsilon, \epsilon'} = \frac{\mathbf{k}}{4} & \text{except } M_{-1,-1} = \frac{\mathbf{k}-4}{4}, \quad \text{for } \frac{\mathbf{k}}{2} \equiv 0 \pmod{2}. \end{cases}$$

Proof. The $\mathbf{f}$ -action (16) on the basis $\mathcal{B}_{\mathbf{c}}$ of $\mathcal{V}_{\mathbf{c}}$ gives rise to the $\langle \mathbf{f} \rangle$ -orbit decomposition,

$$\mathcal{B}_{\mathbf{c}} = \bigsqcup_{\nu=0}^{\lfloor \frac{\mathbf{k}-2}{2} \rfloor} \mathcal{B}_{\mathbf{c}, \nu}, \quad \mathcal{B}_{\mathbf{c}, \nu} := \{[l, m] \in \mathcal{B}_{\mathbf{c}} \mid l = \nu\}.$$

Define $\mathcal{V}_{\mathbf{c}, \nu} = \oplus_{\lambda \in \mathcal{B}_{\mathbf{c}, \nu}} \mathbb{C}\lambda$. As $\Lambda(\mathbf{c})$ -modules, $\mathcal{V}_{\mathbf{c}}$ is the direct sum of $\mathcal{V}_{\mathbf{c}, \nu}$ s, each of which is $\mathbf{k}$ -dimensional except $\dim \mathcal{V}_{\mathbf{c}, \frac{\mathbf{k}-2}{2}} = \frac{\mathbf{k}}{2}$. For odd $\mathbf{k}$, by (18) the $\overline{\Lambda(\mathbf{c})}$ -representation $\mathcal{V}_{\mathbf{c}, \nu}$ is irreducible, and equivalent to $\mathbb{C}_{(1,1)}^N$ for all ν . For even $\mathbf{k}$, the irreducible decomposition of the $\overline{\Lambda(\mathbf{c})}$ -module $\mathcal{V}_{\mathbf{c}, \nu}$ is given by

$$\mathcal{V}_{\mathbf{c}, \nu} = \begin{cases} \mathbb{C}_{(1,1)}^N + \mathbb{C}_{(1,-1)}^N, & \text{if } 0 \leq \nu < \frac{\mathbf{k}-2}{2}, \nu \equiv 0 \pmod{2}, \\ \mathbb{C}_{(-1,1)}^N + \mathbb{C}_{(-1,-1)}^N, & \text{if } 0 \leq \nu < \frac{\mathbf{k}-2}{2}, \nu \equiv 1 \pmod{2}, \\ \mathbb{C}_{(1,1)}^N, & \text{if } \nu = \frac{\mathbf{k}-2}{2}, \frac{\mathbf{k}}{2} \equiv 1 \pmod{2}, \\ \mathbb{C}_{(-1,-1)}^N, & \text{if } \nu = \frac{\mathbf{k}-2}{2}, \frac{\mathbf{k}}{2} \equiv 0 \pmod{2}. \end{cases}$$

Then the result follows immediately. $\square$

For the matrix realization of elliptic and modular transformations, we shall make the identification of vector spaces, $\mathbb{C}^M \otimes \mathbb{C}^N = \mathcal{V}_{\mathbf{c}}$, through a canonical correspondence of the basis elements. For odd $\mathbf{k}$, the identification is made by the following relations for $0 \leq \nu < M, n \in \mathbb{Z}$,

$$|\nu \rangle \otimes |n \rangle = \begin{cases} |\nu, -2n\rangle & \text{for even } \nu, \\ |\nu, \mathbf{k} - 2n\rangle & \text{for odd } \nu. \end{cases} \quad (22)$$

For even $\mathbf{k}$, with $\mathbb{C}^M = \oplus_{\epsilon, \epsilon' = \pm 1} \mathbb{C}^{M, \epsilon, \epsilon'}$ and the standard base of $\mathbb{C}^{M, \epsilon, \epsilon'}$ denoted by $|\nu \rangle_{\epsilon, \epsilon'}$ for $0 \leq \nu < M_{\epsilon, \epsilon'}$, the identification of $\mathbb{C}^M \otimes \mathbb{C}^N$ and $\mathcal{V}_{\mathbf{c}}$ is given by

$$\begin{aligned} |\nu \rangle_{1,1} \otimes |n \rangle &= \frac{1}{2}([2\nu, -2n] + [2\nu, -2n - \mathbf{k}]), & \text{for } 0 \leq \nu < M_{1,1}, \\ |\nu \rangle_{1,-1} \otimes |n \rangle &= \frac{1}{2}([2\nu, -2n] - [2\nu, -2n - \mathbf{k}]), & \text{for } 0 \leq \nu < M_{-1,1}, \\ |\nu \rangle_{-1,1} \otimes |n \rangle &= \frac{1}{2}([2\nu + 1, 1 - 2n] + [2\nu + 1, 1 - 2n - \mathbf{k}]), & \text{for } 0 \leq \nu < M_{-1,1}, \\ |\nu \rangle_{-1,-1} \otimes |n \rangle &= \frac{1}{2}([2\nu + 1, 1 - 2n] - [2\nu + 1, 1 - 2n - \mathbf{k}]), & \text{for } 0 \leq \nu < M_{-1,-1}. \end{aligned} \quad (23)$$

Note that by $\overline{(\frac{\mathbf{k}-2}{2}, m)} = \overline{(\frac{\mathbf{k}-2}{2}, m + \mathbf{k})}$, one has the following identities,

$$\begin{aligned} |\nu \rangle_{1,1} \otimes |n \rangle &= [2\nu, -2n], & \text{for } \nu = M_{1,1} - 1, \frac{\mathbf{k}}{2} \equiv 1 \pmod{2}, \\ |\nu \rangle_{-1,1} \otimes |n \rangle &= [2\nu + 1, 1 - 2n], & \text{for } \nu = M_{-1,1} - 1, \frac{\mathbf{k}}{2} \equiv 0 \pmod{2} \end{aligned}$$

As an example, the identification (23) for $\mathbf{k} = 4$ is given by

$$\begin{aligned} |0 \rangle_{1,1} \otimes |0 \rangle &= \frac{1}{2}[0, 0] + [0, -4], & |0 \rangle_{1,1} \otimes |1 \rangle &= \frac{1}{2}([0, -2] + [0, -6]), \\ |0 \rangle_{1,-1} \otimes |0 \rangle &= \frac{1}{2}([0, 0] - [0, -4]), & |0 \rangle_{1,-1} \otimes |1 \rangle &= \frac{1}{2}([0, -2] - [0, -6]), \\ |0 \rangle_{-1,1} \otimes |0 \rangle &= [1, 1], & |0 \rangle_{-1,1} \otimes |1 \rangle &= [1, -1]. \end{aligned}$$

By (18), one obtains the following results.

Proposition 3. With the identification (22), (23) of $\mathcal{V}_{\mathbf{c}} = \mathbb{C}^M \otimes \mathbb{C}^N$, the $\overline{\Lambda(\mathbf{c})}$ -representation (20) on characters $\text{Ch}_{\lambda}^{(a,b)}(z, \tau)$ has the following matrix form:

- (i) For odd $\mathbf{k}$, $\mathbf{Q} \mapsto \text{Id} \otimes V_N^{-1}$, $\mathbf{P} \mapsto \text{Id} \otimes U_N$.
- (ii) For even $\mathbf{k}$, one has the $\overline{\Lambda(\mathbf{c})}$ -decomposition $\mathbb{C}^M \otimes \mathbb{C}^N = \oplus_{\epsilon, \epsilon' = \pm 1} (\mathbb{C}^{M_{\epsilon, \epsilon'}} \otimes \mathbb{C}^N)$ with the matrix expressions on $\mathbb{C}^{M_{\epsilon, \epsilon'}} \otimes \mathbb{C}^N$ given by

$$\begin{aligned} \text{on } \mathbb{C}^{M_{1, \epsilon'}} \otimes \mathbb{C}^N : \quad & \mathbf{Q} \mapsto \text{Id} \otimes V_N^{-1}, \quad \mathbf{P} \mapsto \text{Id} \otimes U_N^{\epsilon'}, \\ \text{on } \mathbb{C}^{M_{-1, \epsilon'}} \otimes \mathbb{C}^N : \quad & \mathbf{Q} \mapsto \omega^{\frac{1}{2}} \text{Id} \otimes V_N^{-1}, \quad \mathbf{P} \mapsto \text{Id} \otimes U_N^{\epsilon'}, \end{aligned}$$

where $\epsilon' = \pm 1$. $\square$

For the discussion of the modular properties of the characters $\text{Ch}_{\lambda}^{(a,b)}(z, \tau)$, the modular group is Γ_q with $q = (\frac{1}{2} - a, \frac{1}{2} - b)$. Its description is given by (5) with $d = \mathbf{c}, w = 0$ while restricting on Γ_q . Together with the elliptic action (20), the relations among characters $\text{Ch}_{\lambda}^{(a,b)}(z, \tau)$ s give rise to a finite dimensional representation of a group $\overline{\Lambda(\mathbf{c})} \overset{r}{*} \Gamma_q$ through (7) for $r = (a, b), d = \mathbf{c}, w = 0$. Here the group $\overline{\Lambda(\mathbf{c})} \overset{r}{*} \Gamma_q$, is defined by the following exact sequence using the Γ_q -conjugation invariance property of the subgroup $\mathbf{k}\Lambda$ of $\Lambda(\mathbf{c})$,

$$0 \longrightarrow \mathbf{k}\Lambda \overset{r}{*} \Gamma_q \longrightarrow \Lambda(\mathbf{c}) \overset{r}{*} \Gamma_q \longrightarrow \overline{\Lambda(\mathbf{c})} \overset{r}{*} \Gamma_q \longrightarrow 0.$$

In particular, Γ_q is the full modular group Γ for the Ramon characters R_{λ} or $\tilde{R}_{\lambda}$; $\Gamma_{(\frac{1}{2}, \frac{1}{2})}$ for the Neveu-Schwarz characters NS_{λ} . In this paper, we shall only consider these two special cases, partly due to the easy description of generators of these modular groups, but also its sufficiency for the future study on topological applications of manifolds. We denote

$$\Gamma_{(\frac{1}{2}, \frac{1}{2})}^{\text{W}}(\mathbf{c}) := \overline{\Lambda(\mathbf{c})} \overset{r}{*} \Gamma_{(\frac{1}{2}, \frac{1}{2})}, \quad \Gamma^{\text{W}}(\mathbf{c}) := \overline{\Lambda(\mathbf{c})} \overset{(\frac{1}{2}, \frac{1}{2})}{*} \Gamma.$$

We now describe the T, S-transforms on Ramon and Neveu-Schwarz characters.

Proposition 4. The following relations hold:

$$\begin{aligned} \text{NS}_{\lambda}(z, \tau + 2) &= e^{4\pi\sqrt{-1}(H_{\lambda} - \frac{\mathbf{s}}{8})} \text{NS}_{\lambda}(z, \tau), \quad e^{\frac{-c\pi\sqrt{-1}z^2}{\tau}} \text{NS}_{\lambda}(\frac{z}{-\tau}, \frac{1}{-\tau}) = \sum_{\lambda' \in \mathcal{B}_{\mathbf{c}}} S_{\lambda}^{\lambda'} \text{NS}_{\lambda'}(z, \tau), \\ R_{\lambda}(z, \tau + 1) &= e^{2\pi\sqrt{-1}(H_{\lambda} + \frac{Q_{\lambda}}{2})} R_{\lambda}(z, \tau), \quad e^{\frac{-c\pi\sqrt{-1}z^2}{\tau}} R_{\lambda}(\frac{z}{-\tau}, \frac{1}{-\tau}) = e^{2\pi\sqrt{-1}(Q_{\lambda} + \frac{\mathbf{s}}{4})} \sum_{\lambda' \in \mathcal{B}_{\mathbf{c}}} S_{\lambda}^{\lambda'} R_{\lambda'}(z, \tau), \\ \tilde{R}_{\lambda}(z, \tau + 1) &= e^{2\pi\sqrt{-1}(H_{\lambda} - \frac{Q_{\lambda}}{2})} \tilde{R}_{\lambda}(z, \tau), \quad e^{\frac{-c\pi\sqrt{-1}z^2}{\tau}} \tilde{R}_{\lambda}(\frac{z}{-\tau}, \frac{1}{-\tau}) = e^{2\pi\sqrt{-1}(-Q_{\lambda} + \frac{\mathbf{s}}{4})} \sum_{\lambda' \in \mathcal{B}_{\mathbf{c}}} S_{\lambda}^{\lambda'} \tilde{R}_{\lambda'}(z, \tau), \end{aligned}$$

where the coefficients $S_{\lambda}^{\lambda'}$, $\lambda = [l, m], \lambda' = [l', m']$, are defined by

$$S_{\lambda}^{\lambda'} := \frac{2}{\mathbf{k}} \sin \frac{\pi(l+1)(l'+1)}{\mathbf{k}} e^{\frac{-\pi\sqrt{-1}mm'}{\mathbf{k}}} = \frac{2}{\mathbf{k}} \sin \frac{\pi(i-j)(i'-j')}{\mathbf{k}} e^{\frac{-\pi\sqrt{-1}(i+j)(i'+j')}{\mathbf{k}}}.$$

Proof. The T-relation of $R_{\lambda}, \tilde{R}_{\lambda}$ follows from (15) (9). By (10) for $(a, b) = (\frac{1}{2}, \frac{1}{2})$ and the T-relation of Ramon characters, one obtains the T²-relation of NS_{λ} . The S-transform of NS_{λ} was known in literature, (e.g., formula (28a) of [17]), which we adopt here. Then, by (10), we have the relation,

$$e^{\frac{-c\pi\sqrt{-1}z^2}{\tau}} \text{Ch}_{\lambda}^{(a,b)}(\frac{z}{-\tau}, \frac{1}{-\tau}) = e^{2abc\pi\sqrt{-1}} \sum_{\lambda' \in \mathcal{B}_{\mathbf{c}}} S_{\lambda}^{\lambda'} \text{Ch}_{\lambda'}^{(-b,a)}(z, \tau).$$

Setting $(a, b) = (\frac{1}{2}, \frac{-1}{2})$ or $(\frac{-1}{2}, \frac{1}{2})$ in the above equality, one obtains the expression of the S-transform of Ramon characters by (11) (19). $\square$

With the identification of the basis element λ of $\mathcal{V}_{\mathbf{c}}$ with the character NS_{λ} or $\text{R}_{\lambda}, \tilde{\text{R}}_{\lambda}$, by (20), by using Proposition 4 and the remark of Proposition 2, there exists a canonical Jacobi-group representation on $\mathcal{V}_{\mathbf{c}}$ for the Neveu-Schwarz or Ramon sector,

$$\Pi_{\text{NS}} : \mathcal{V}_{\mathbf{c}} \times \Gamma_{(\frac{1}{2}, \frac{1}{2})}^{\text{W}}(\mathbf{c}) \longrightarrow \mathcal{V}_{\mathbf{c}}, \quad \Pi_{\text{R}}, \Pi_{\tilde{\text{R}}} : \mathcal{V}_{\mathbf{c}} \times \Gamma^{\text{W}}(\mathbf{c}) \longrightarrow \mathcal{V}_{\mathbf{c}}.$$

With $\mathcal{V}_{\mathbf{c}} = \mathbb{C}^M \otimes \mathbb{C}^N$ through (22) (23), the matrix presentation of the above actions on the part $\overline{\Lambda(\mathbf{c})}$ is described by Proposition 3. As any modular transformation sends a $\overline{\Lambda(\mathbf{c})}$ -irreducible subspace onto another one, we shall be able to obtain a matrix form of a modular transformation as a tensor product of one from the $\mathbb{C}^M$ -factor, $(\alpha_{\nu\nu'})_{\nu, \nu'}$, and another one from the $\mathbb{C}^N$ -factor, $(\beta_{nn'})_{n, n'}$. By Proposition 4, T-transformation is a diagonal matrix; while the S-transform of a vector will be a linear combination of all the basis elements. The explicit forms are given in the following theorem, where the matrix indices ν, ν', n, n' are non-negative integers less than the dimensions of vector spaces involved in the expression.

Theorem 2. For $\Pi_{\text{NS}}, \Pi_{\text{R}}, \Pi_{\tilde{\text{R}}}$ -representations, the matrix forms of the modular transformations are as follows:

(i) For odd $\mathbf{k}$, with $\mathcal{V}_{\mathbf{c}} = \mathbb{C}^M \otimes \mathbb{C}^N$ by (22), we have

$$\begin{aligned} \Pi_{\text{NS}} : \quad \text{T}^2 &\mapsto -\sqrt{-1}((-1)^{\frac{(\nu+1)^{\nu}-1}{2}} e^{\frac{(\nu+1)^2 \pi \sqrt{-1}}{\mathbf{k}}} \delta_{\nu, \nu'}) \otimes (\omega^{-n^2} \delta_{n, n'}), \\ \text{S} &\mapsto \frac{2}{\mathbf{k}} (\sin \frac{(\nu+1)(\nu'+1)\pi}{\mathbf{k}}) \otimes (\omega^{-nn'}), \\ \Pi_{\text{R}} : \quad \text{T} &\mapsto ((-1)^{\frac{(\nu+1)^{\nu}-1}{4}} e^{\frac{(\nu+1)^2 \pi \sqrt{-1}}{2\mathbf{k}}} \delta_{\nu, \nu'}) \otimes (\omega^{\frac{-(2n+1)^2}{8}} \delta_{n, n'}), \\ \text{S} &\mapsto \frac{2\sqrt{-1}}{\mathbf{k}} (\sin \frac{(\nu+1)(\nu'+1)\pi}{\mathbf{k}}) \otimes (\omega^{-n(n'+1)-\frac{1}{4}}), \\ \Pi_{\tilde{\text{R}}} : \quad \text{T} &\mapsto ((-1)^{\frac{(\nu+1)^{\nu}-1}{4}} e^{\frac{(\nu+1)^2 \pi \sqrt{-1}}{2\mathbf{k}}} \delta_{\nu, \nu'}) \otimes (\omega^{\frac{-(2n-1)^2}{8}} \delta_{n, n'}), \\ \text{S} &\mapsto \frac{2\sqrt{-1}}{\mathbf{k}} (\sin \frac{(\nu+1)(\nu'+1)\pi}{\mathbf{k}}) \otimes (\omega^{-n(n'-1)-\frac{1}{4}}), \end{aligned}$$

where $(-1)^* := e^{*(\mathbf{k}-2)\pi\sqrt{-1}}$.

(ii) For even $\mathbf{k}$, with $\mathcal{V}_{\mathbf{c}} = \oplus_{\epsilon, \epsilon'=\pm 1, 2} \mathbb{C}^{M_{\epsilon, \epsilon'}} \otimes \mathbb{C}^N$ by (23), the linear transformations, T, S, are composed of the following ones on the Neveu-Schwarz, or Ramon sector:

$$\begin{aligned} \Pi_{\text{NS}}, \quad \text{T}^2 : \mathbb{C}^{M_{\epsilon, \epsilon'}} \otimes \mathbb{C}^N &\longrightarrow \mathbb{C}^{M_{\epsilon, \epsilon'}} \otimes \mathbb{C}^N, \\ \Pi_{\text{R}}, \Pi_{\tilde{\text{R}}}, \quad \text{T} : \mathbb{C}^{M_{\epsilon, \epsilon'}} \otimes \mathbb{C}^N &\longrightarrow \mathbb{C}^{M_{\epsilon, \epsilon''}} \otimes \mathbb{C}^N, \\ \Pi_{\text{NS}}, \Pi_{\text{R}}, \Pi_{\tilde{\text{R}}}, \quad \text{S} : \mathbb{C}^{M_{\epsilon, \epsilon'}} \otimes \mathbb{C}^N &\longrightarrow \mathbb{C}^{M_{\epsilon', \epsilon}} \otimes \mathbb{C}^N, \end{aligned}$$

where $\epsilon, \epsilon' = \pm 1$, and $\epsilon'' := (-1)^{\frac{\mathbf{k}-2}{2}} \epsilon \epsilon'$. The matrix representation of the above linear transformations is given by

$$\begin{aligned} \Pi_{\text{NS}}, \quad \mathbb{C}^{M_{\epsilon, \epsilon'}} \otimes \mathbb{C}^N &\xrightarrow{\text{T}^2} \mathbb{C}^{M_{\epsilon, \epsilon'}} \otimes \mathbb{C}^N, \quad \mapsto -\sqrt{-1} (e^{\frac{(4\nu+3-\epsilon)^2 \pi \sqrt{-1}}{4\mathbf{k}}} \delta_{\nu, \nu'}) \otimes (\omega^{\frac{-(4n-1+\epsilon)^2}{16}} \delta_{n, n'}), \\ \Pi_{\text{R}}, \quad \mathbb{C}^{M_{\epsilon, \epsilon'}} \otimes \mathbb{C}^N &\xrightarrow{\text{T}} \mathbb{C}^{M_{\epsilon, \epsilon''}} \otimes \mathbb{C}^N, \quad \mapsto (e^{\frac{(4\nu+3-\epsilon)^2 \pi \sqrt{-1}}{8\mathbf{k}}} \delta_{\nu, \nu'}) \otimes (\omega^{\frac{-(4n+1+\epsilon)^2}{32}} \delta_{n, n'}), \\ \Pi_{\tilde{\text{R}}}, \quad \mathbb{C}^{M_{\epsilon, \epsilon'}} \otimes \mathbb{C}^N &\xrightarrow{\text{T}} \mathbb{C}^{M_{\epsilon, \epsilon''}} \otimes \mathbb{C}^N, \quad \mapsto (e^{\frac{(4\nu+3-\epsilon)^2 \pi \sqrt{-1}}{8\mathbf{k}}} \delta_{\nu, \nu'}) \otimes (\omega^{\frac{-(4n-3+\epsilon)^2}{32}} \delta_{n, n'}), \end{aligned}$$

and

$$\begin{aligned}
\mathbb{C}^{M_{1,1}} \otimes \mathbb{C}^N &\xrightarrow{S} \mathbb{C}^{M_{1,1}} \otimes \mathbb{C}^N, & \mapsto \begin{cases} \frac{4}{k}(w_{1,1}(\nu') \sin \frac{\pi(2\nu+1)(2\nu'+1)}{k}) \otimes (\omega^{nn'}) & \text{for } \Pi_{\text{NS}}, \\ \frac{4\sqrt{-1}}{k}(w_{1,1}(\nu') \sin \frac{\pi(2\nu+1)(2\nu'+1)}{k}) \otimes (\omega^{n(n'-1)-\frac{1}{4}}) & \text{for } \Pi_{\text{R}}, \\ \frac{4\sqrt{-1}}{k}(w_{1,1}(\nu') \sin \frac{\pi(2\nu+1)(2\nu'+1)}{k}) \otimes (\omega^{n(n'+1)-\frac{1}{4}}) & \text{for } \Pi_{\tilde{\text{R}}}, \end{cases} \\
\mathbb{C}^{M_{-1,1}} \otimes \mathbb{C}^N &\xrightarrow{S} \mathbb{C}^{M_{-1,1}} \otimes \mathbb{C}^N, & \mapsto \begin{cases} \frac{4}{k}(w_{-1,1}(\nu') \sin \frac{\pi(2\nu+1)(2\nu'+2)}{k}) \otimes (\omega^{n(n'-\frac{1}{2})}) & \text{for } \Pi_{\text{NS}}, \\ \frac{4\sqrt{-1}}{k}(w_{-1,1}(\nu') \sin \frac{\pi(2\nu+1)(2\nu'+2)}{k}) \otimes (\omega^{n(n'-\frac{3}{2})-\frac{1}{4}}) & \text{for } \Pi_{\text{R}}, \\ \frac{4\sqrt{-1}}{k}(w_{-1,1}(\nu') \sin \frac{\pi(2\nu+1)(2\nu'+2)}{k}) \otimes (\omega^{n(n'+\frac{1}{2})-\frac{1}{4}}) & \text{for } \Pi_{\tilde{\text{R}}}, \end{cases} \\
\mathbb{C}^{M_{-1,1}} \otimes \mathbb{C}^N &\xrightarrow{S} \mathbb{C}^{M_{1,-1}} \otimes \mathbb{C}^N, & \mapsto \begin{cases} \frac{4}{k}(\sin \frac{\pi(2\nu+2)(2\nu'+1)}{k}) \otimes (\omega^{(n-\frac{1}{2})n'}) & \text{for } \Pi_{\text{NS}}, \\ \frac{4\sqrt{-1}}{k}(\sin \frac{\pi(2\nu+2)(2\nu'+1)}{k}) \otimes (\omega^{(n-\frac{1}{2})n'-n+\frac{1}{4}}) & \text{for } \Pi_{\text{R}}, \\ \frac{4\sqrt{-1}}{k}(\sin \frac{\pi(2\nu+2)(2\nu'+1)}{k}) \otimes (\omega^{(n-\frac{1}{2})n'+n-\frac{3}{4}}) & \text{for } \Pi_{\tilde{\text{R}}}, \end{cases} \\
\mathbb{C}^{M_{-1,-1}} \otimes \mathbb{C}^N &\xrightarrow{S} \mathbb{C}^{M_{-1,-1}} \otimes \mathbb{C}^N, & \mapsto \begin{cases} \frac{4}{k}(\sin \frac{\pi(2\nu+2)(2\nu'+2)}{k}) \otimes (\omega^{(n-\frac{1}{2})(n'-\frac{1}{2})}) & \text{for } \Pi_{\text{NS}}, \\ \frac{4\sqrt{-1}}{k}(\sin \frac{\pi(2\nu+2)(2\nu'+2)}{k}) \otimes (\omega^{(n-\frac{1}{2})(n'-\frac{1}{2})-n+\frac{1}{4}}) & \text{for } \Pi_{\text{R}}, \\ \frac{4\sqrt{-1}}{k}(\sin \frac{\pi(2\nu+2)(2\nu'+2)}{k}) \otimes (\omega^{(n-\frac{1}{2})(n'-\frac{1}{2})+n-\frac{3}{4}}) & \text{for } \Pi_{\tilde{\text{R}}}. \end{cases}
\end{aligned}$$

Here the functions, $w_{1,1}(\nu')$, $w_{-1,1}(\nu')$, are defined by

$$\begin{aligned}
w_{1,1}(\nu') &= \begin{cases} \frac{1}{2} & \text{if } \nu' = M_{1,1} - 1, \frac{k}{2} \equiv 1 \pmod{2}, \\ 1 & \text{otherwise with } 0 \leq \nu' < M_{1,1}, \end{cases} \\
w_{-1,1}(\nu') &= \begin{cases} \frac{1}{2} & \text{if } \nu' = M_{-1,1} - 1, \frac{k}{2} \equiv 0 \pmod{2}, \\ 1 & \text{otherwise with } 0 \leq \nu' < M_{-1,1}. \end{cases}
\end{aligned}$$

Proof. (i) For odd k , we have $k = N = 2M + 1$. By Proposition 4 and (12), one has the following expressions of the T-transformation on the Neveu-Schwarz or Ramon sector,

$$\begin{aligned}
\Pi_{\text{NS}} : \quad & T^2(|\nu > \otimes |n >) = -\sqrt{-1}(-1)^{\frac{(\nu+1)^2-1}{2}} e^{\frac{(\nu+1)^2\pi\sqrt{-1}}{k}} \omega^{-n^2} |\nu > \otimes |n >, \\
& S(|\nu > \otimes |n >) = \frac{2}{k} \sum_{\nu'=0}^{M-1} \sum_{n'=0}^{N-1} \sin \frac{(\nu+1)(\nu'+1)\pi}{k} \omega^{-nn'} |\nu' > \otimes |n' >, \\
\Pi_{\text{R}} : \quad & T(|\nu > \otimes |n >) = (-1)^{\frac{(\nu+1)^2-1}{4}} e^{\frac{(\nu+1)^2\pi\sqrt{-1}}{2k}} \omega^{-\frac{(2n+1)^2}{8}} |\nu > \otimes |n >, \\
& S(|\nu > \otimes |n >) = \frac{2\sqrt{-1}}{k} \sum_{\nu'=0}^{M-1} \sum_{n'=0}^{N-1} \sin \frac{(\nu+1)(\nu'+1)\pi}{k} \omega^{-n(n'+1)-\frac{1}{4}} |\nu' > \otimes |n' >, \\
\Pi_{\tilde{\text{R}}} : \quad & T(|\nu > \otimes |n >) = (-1)^{\frac{(\nu+1)^2-1}{4}} e^{\frac{(\nu+1)^2\pi\sqrt{-1}}{2k}} \omega^{-\frac{(2n-1)^2}{8}} |\nu > \otimes |n >, \\
& S(|\nu > \otimes |n >) = \frac{2\sqrt{-1}}{k} \sum_{\nu'=0}^{M-1} \sum_{n'=0}^{N-1} \sin \frac{(\nu+1)(\nu'+1)\pi}{k} \omega^{-n(n'-1)-\frac{1}{4}} |\nu' > \otimes |n' >.
\end{aligned}$$

Then the results follow immediately.

(ii) When k is even, we have $k = 2N = M + 1$, and $\omega^{\frac{k}{2}} = 1$. One has the following identities for $\lambda = [l, m]$, $\lambda' = [l, m - k]$,

$$e^{4\pi\sqrt{-1}(H_{\lambda'})} = e^{4\pi\sqrt{-1}(H_{\lambda})}, \quad e^{2\pi\sqrt{-1}(H_{\lambda'} \pm \frac{Q_{\lambda'}}{2})} = (-1)^{\frac{k-2}{2}+m} e^{2\pi\sqrt{-1}(H_{\lambda} \pm \frac{Q_{\lambda}}{2})}, \quad e^{2\pi\sqrt{-1}Q_{\lambda'}} = e^{2\pi\sqrt{-1}Q_{\lambda}}.$$

By the first two equalities in the above and Proposition 4, one obtains the statement on T^2 or T on the $\mathbb{C}^{M_{\epsilon\epsilon'}} \otimes \mathbb{C}^N$ for the Neveu-Schwarz, Ramon sector respectively. Furthermore, by

$$\begin{aligned}
e^{4\pi\sqrt{-1}(H_{[2\nu,-2n]} - \frac{\epsilon}{8})} &= -\sqrt{-1} e^{\frac{(2\nu+1)^2\pi\sqrt{-1}}{k}} \omega^{-n^2}, \quad e^{4\pi\sqrt{-1}(H_{[2\nu+1,1-2n]} - \frac{\epsilon}{8})} = -\sqrt{-1} e^{\frac{(2\nu+2)^2\pi\sqrt{-1}}{k}} \omega^{-(n-\frac{1}{2})^2}, \\
e^{2\pi\sqrt{-1}(H_{[2\nu,-2n]} \pm \frac{Q_{[2\nu,-2n]}}{2})} &= e^{\frac{(2\nu+1)^2\pi\sqrt{-1}}{2k}} \omega^{-\frac{(n\pm 1/2)^2}{2}}, \\
e^{2\pi\sqrt{-1}(H_{[2\nu+1,1-2n]} \pm \frac{Q_{[2\nu+1,1-2n]}}{2})} &= e^{\frac{(2\nu+2)^2\pi\sqrt{-1}}{2k}} \omega^{-\frac{(n-\frac{1}{2} \pm \frac{1}{2})^2}{2}},
\end{aligned}$$

the expressions of T-transformation on the basis elements of $\mathbb{C}^{M_{\epsilon\epsilon'}} \otimes \mathbb{C}^N$ are as follows,

$$\begin{aligned}
\Pi_{\text{NS}} : \quad & T^2(|\nu >_{1,\epsilon'} \otimes |n >) = -\sqrt{-1} e^{\frac{(2\nu+1)^2 \pi \sqrt{-1}}{\mathbf{k}}} \omega^{-n^2} |\nu >_{1,\epsilon'} \otimes |n >, \\
& T^2(|\nu >_{-1,\epsilon'} \otimes |n >) = -\sqrt{-1} e^{\frac{(2\nu+2)^2 \pi \sqrt{-1}}{\mathbf{k}}} \omega^{-(n-\frac{1}{2})^2} |\nu >_{-1,\epsilon'} \otimes |n >, \\
\Pi_{\text{R}} : \quad & T(|\nu >_{1,\epsilon'} \otimes |n >) = e^{\frac{(2\nu+1)^2 \pi \sqrt{-1}}{2\mathbf{k}}} \omega^{\frac{-(n+1/2)^2}{2}} |\nu >_{1,\epsilon''} \otimes |n >, \\
& T(|\nu >_{-1,\epsilon'} \otimes |n >) = e^{\frac{(2\nu+2)^2 \pi \sqrt{-1}}{2\mathbf{k}}} \omega^{\frac{-n^2}{2}} |\nu >_{-1,\epsilon''} \otimes |n >, \\
\Pi_{\widetilde{\text{R}}} : \quad & T(|\nu >_{1,\epsilon'} \otimes |n >) = e^{\frac{(2\nu+1)^2 \pi \sqrt{-1}}{2\mathbf{k}}} \omega^{\frac{-(n-1/2)^2}{2}} |\nu >_{1,\epsilon''} \otimes |n >, \\
& T(|\nu >_{-1,\epsilon'} \otimes |n >) = e^{\frac{(2\nu+2)^2 \pi \sqrt{-1}}{2\mathbf{k}}} \omega^{\frac{-(n-1)^2}{2}} |\nu >_{-1,\epsilon''} \otimes |n >.
\end{aligned}$$

Hence we obtain the matrix expression of T^2 for Π_{NS} ; and T for $\Pi_{\text{R}}, \Pi_{\widetilde{\text{R}}}$ -action. For the S-transformation, we first consider the Neveu-Schwarz sector. By the definition of $S_{\lambda}^{\lambda'}$, the following equalities hold: for even ν ,

$$\begin{aligned}
S([\nu, -2n]) &= \frac{2}{\mathbf{k}} \sum_{\text{even } \nu'=0}^{\frac{\mathbf{k}-2}{2}-1} \sin \frac{\pi(\nu+1)(\nu'+1)}{\mathbf{k}} \sum_{n'=0}^{\frac{\mathbf{k}}{2}-1} \omega^{nn'} ([\nu', -2n'] + [\nu', -2n' - \mathbf{k}]) \\
&+ \frac{2}{\mathbf{k}} \sum_{\text{odd } \nu'=0}^{\frac{\mathbf{k}-2}{2}-1} \sin \frac{\pi(\nu+1)(\nu'+1)}{\mathbf{k}} \sum_{n'=0}^{\frac{\mathbf{k}}{2}-1} \omega^{n(n'-\frac{1}{2})} ([\nu', 1 - 2n'] + [\nu', 1 - 2n' - \mathbf{k}]) \\
&+ \frac{2}{\mathbf{k}} \sin \frac{\pi(\nu+1)}{2} \sum_{n'=0}^{\frac{\mathbf{k}}{2}-1} \omega^{n(n'-\frac{\theta(\mathbf{k})}{2})} [\frac{\mathbf{k}-2}{2}, \theta(\mathbf{k}) - 2n'] , \\
S([\nu, -2n - \mathbf{k}]) &= \frac{2}{\mathbf{k}} \sum_{\text{even } \nu'=0}^{\frac{\mathbf{k}-2}{2}-1} \sin \frac{\pi(\nu+1)(\nu'+1)}{\mathbf{k}} \sum_{n'=0}^{\frac{\mathbf{k}}{2}-1} \omega^{nn'} ([\nu', -2n'] + [\nu', -2n' - \mathbf{k}]) \\
&- \frac{2}{\mathbf{k}} \sum_{\text{odd } \nu'=0}^{\frac{\mathbf{k}-2}{2}-1} \sin \frac{\pi(\nu+1)(\nu'+1)}{\mathbf{k}} \sum_{n'=0}^{\frac{\mathbf{k}}{2}-1} \omega^{n(n'-\frac{1}{2})} ([\nu', 1 - 2n'] + [\nu', 1 - 2n' - \mathbf{k}]) \\
&+ \frac{2}{\mathbf{k}} \sin \frac{\pi(\nu+1)}{2} \sum_{n'=0}^{\frac{\mathbf{k}}{2}-1} \omega^{n(n'-\frac{\theta(\mathbf{k})}{2})} [\frac{\mathbf{k}-2}{2}, \theta(\mathbf{k}) - 2n'] ,
\end{aligned}$$

where $\theta(\mathbf{k}) := 0, 1$ according to $\frac{\mathbf{k}}{2}$ odd or even respectively ; for odd ν ,

$$\begin{aligned}
S([\nu, 1 - 2n]) &= \frac{2}{\mathbf{k}} \sum_{\text{even } \nu'=0}^{\frac{\mathbf{k}-2}{2}-1} \sin \frac{\pi(\nu+1)(\nu'+1)}{\mathbf{k}} \sum_{n'=0}^{\frac{\mathbf{k}}{2}-1} \omega^{(n-\frac{1}{2})n'} ([\nu', -2n'] - [\nu', -2n' - \mathbf{k}]) \\
&+ \frac{2}{\mathbf{k}} \sum_{\text{odd } \nu'=0}^{\frac{\mathbf{k}-2}{2}-1} \sin \frac{\pi(\nu+1)(\nu'+1)}{\mathbf{k}} \sum_{n'=0}^{\frac{\mathbf{k}}{2}-1} \omega^{(n-\frac{1}{2})(n'-\frac{1}{2})} ([\nu', 1 - 2n'] - [\nu', 1 - 2n' - \mathbf{k}]) , \\
S([\nu, 1 - 2n - \mathbf{k}]) &= \frac{2}{\mathbf{k}} \sum_{\text{even } \nu'=0}^{\frac{\mathbf{k}-2}{2}-1} \sin \frac{\pi(\nu+1)(\nu'+1)}{\mathbf{k}} \sum_{n'=0}^{\frac{\mathbf{k}}{2}-1} \omega^{(n-\frac{1}{2})n'} ([\nu', -2n'] - [\nu', -2n' - \mathbf{k}]) \\
&- \frac{2}{\mathbf{k}} \sum_{\text{odd } \nu'=0}^{\frac{\mathbf{k}-2}{2}-1} \sin \frac{\pi(\nu+1)(\nu'+1)}{\mathbf{k}} \sum_{n'=0}^{\frac{\mathbf{k}}{2}-1} \omega^{(n-\frac{1}{2})(n'-\frac{1}{2})} ([\nu', 1 - 2n'] - [\nu', 1 - 2n' - \mathbf{k}]) .
\end{aligned}$$

From which one can derive the following relations,

$$\begin{aligned}
S(|\nu >_{1,1} \otimes |n >) &= \frac{4}{\mathbf{k}} \sum_{\nu'=0}^{M_{1,1}-1} \sum_{n'=0}^{N-1} w_{1,1}(\nu') \sin \frac{\pi(2\nu+1)(2\nu'+1)}{\mathbf{k}} \omega^{nn'} |\nu' >_{1,1} \otimes |n' >, \\
S(|\nu >_{1,-1} \otimes |n >) &= \frac{4}{\mathbf{k}} \sum_{\nu'=0}^{M_{-1,1}-1} \sum_{n'=0}^{N-1} w_{-1,1}(\nu') \sin \frac{\pi(2\nu+1)(2\nu'+2)}{\mathbf{k}} \omega^{n(n'-\frac{1}{2})} |\nu' >_{-1,1} \otimes |n' >, \\
S(|\nu >_{-1,1} \otimes |n >) &= \frac{4}{\mathbf{k}} (\sum_{\nu'=0}^{M_{1,-1}-1} \sum_{n'=0}^{N-1} \sin \frac{\pi(2\nu+2)(2\nu'+1)}{\mathbf{k}} \omega^{(n-\frac{1}{2})n'} |\nu' >_{1,-1} \otimes |n' >, \\
S(|\nu >_{-1,-1} \otimes |n >) &= \frac{4}{\mathbf{k}} \sum_{\nu'=0}^{M_{-1,-1}-1} \sum_{n'=0}^{N-1} \sin \frac{\pi(2\nu+2)(2\nu'+2)}{\mathbf{k}} \omega^{(n-\frac{1}{2})(n'-\frac{1}{2})} |\nu' >_{-1,-1} \otimes |n' >.
\end{aligned}$$

Hence we obtain the matrices of S for Π_{NS} . By Proposition 4 the S-expression of a Ramon sector differs from the one of the Neveu-Schwarz sector only by a factor $e^{2\pi\sqrt{-1}(\pm Q_{[2\nu+1,1-2n]} + \frac{\mathbf{e}}{4})}$. Therefore, with the equalities,

$$\begin{aligned}
e^{2\pi\sqrt{-1}(\pm Q_{[2\nu,-2n]} + \frac{\mathbf{e}}{4})} &= \sqrt{-1} \omega^{\mp n-1/4}, \quad e^{2\pi\sqrt{-1}(Q_{[2\nu+1,1-2n]} + \frac{\mathbf{e}}{4})} = \sqrt{-1} \omega^{-n+1/4}, \\
e^{2\pi\sqrt{-1}(-Q_{[2\nu+1,1-2n]} + \frac{\mathbf{e}}{4})} &= \sqrt{-1} \omega^{n-\frac{3}{4}}.
\end{aligned}$$

the S-transformation matrices for $\Pi_{\text{R}}, \Pi_{\widetilde{\text{R}}}$ follow from that of Π_{NS} . $\square$

5 Acknowledgements

I would like to thank MSRI and Mathematical Department of U.C. Berkeley, U. S. A. for the hospitality during the academic year of 1997, where this work was formed. Part of this work was performed during a visit of the author to the Hong Kong University of Science and Technology (HKUST), Hong Kong on November, 1998. I thank Professor Xiaoping Xu for his kind invitation and hospitality at HKUST.

6 Appendix: Geometrical Realization of N=2 Conformal Algebra

The geometrical representation of generators of Virasoro algebra in terms of vector fields on $\mathbb{C}^*$ has been well-known in literature, so is the N=1 conformal algebra's generators as derivations of $\mathbb{C}^*$ with one Grassmann variable θ , (see for instance [2]). The super-vector field expressions of N-extended conformal algebra for a general N have been treated in a similar way, and its various forms could be found in literature [10]. In this section, we are going to present a detailed geometrical realization of the N= 2 conformal algebra so that one could have a better understanding of the symmetries represented by this algebra. We start with an alternative definition of the N=2 conformal algebra, whose formulation works also for an arbitrary value N,

$$\begin{aligned} [L_m, L_n] &= (m-n)L_{m+n} + \frac{c}{12}(m^3-m)\delta_{m+n,0}, & [J_m, J_n] &= \frac{c}{3}m\delta_{m+n,0}, \\ [L_m, J_n] &= -nJ_{m+n}, \quad [L_m, G_p^j] = (\frac{m}{2}-p)G_{m+p}^j, & [J_m, G_p^j] &= \sqrt{-1}\epsilon_{j,k}G_{m+p}^k, \\ \{G_p^j, G_q^k\} &= 2\delta_{j,k}L_{p+q} + \sqrt{-1}\epsilon_{j,k}(p-q)J_{p+q} + \frac{c}{3}(p^2 - \frac{1}{4})\delta_{j,k}\delta_{p+q,0}, \end{aligned}$$

where $j, k = 1, 2$, and $\epsilon_{j,k}$ are antisymmetric with $\epsilon_{1,2} = 1$. The definition of SCA in Sect. 3 is equivalent to the above one by the relations,

$$G_n^+ = \frac{1}{\sqrt{2}}(G_n^1 + \sqrt{-1}G_n^2), \quad G_n^- = \frac{1}{\sqrt{2}}(G_n^1 - \sqrt{-1}G_n^2).$$

It is easy to see that the correspondence,

$$L_n \mapsto (-1)^{n+1}L_{-n}, \quad J_n \mapsto (-1)^n J_{-n}, \quad G_p^* \mapsto (-1)^{p-\frac{1}{2}}G_{-p}^*, \quad (* = 1, 2, \text{ or } \pm), \quad (24)$$

together with $c \mapsto -c$, give rise to an automorphism of the superalgebra SCA. There is a finite dimensional Lie-superalgebra $\mathfrak{g}$ invariant under the above automorphism, the even part $\mathfrak{g}^e$ and odd part $\mathfrak{g}^o$ are given by

$$\begin{aligned} \mathfrak{g}^e &= \langle L_{-1}, L_0, L_1, J_0 \rangle_{\mathbb{C}} \simeq sl_2 \oplus \mathbb{C}J_0, \\ \mathfrak{g}^o &= \langle G_{\frac{\pm 1}{2}}^1, G_{\frac{\pm 1}{2}}^2 \rangle_{\mathbb{C}} = \langle G_{\frac{1}{2}}^{\pm}, G_{\frac{-1}{2}}^{\pm} \rangle_{\mathbb{C}}. \end{aligned}$$

Here the standard basis elements e, f, h of sl_2 with $[h, e] = 2e, [h, f] = -2f, [e, f] = h$, are expressed by

$$e = L_1, \quad h = -2L_0, \quad f = -L_{-1}.$$

The representation of the even algebra $\mathfrak{g}^e$ on the odd space $\mathfrak{g}^o$ decomposes into two copies of the canonical 2-dimensional representation of sl_2 :

$$[h, G_{\frac{\pm 1}{2}}^*] = \pm G_{\frac{\pm 1}{2}}^*, \quad [e, G_{\frac{1}{2}}^*] = 0, \quad [e, G_{\frac{-1}{2}}^*] = G_{\frac{1}{2}}^*, \quad [f, G_{\frac{1}{2}}^*] = G_{\frac{-1}{2}}^*, \quad [f, G_{\frac{-1}{2}}^*] = 0,$$

for $* = 1, 2$, or $\pm$; while the action of J_0 is given by

$$[J_0, G_p^j] = \sqrt{-1}\epsilon_{j,k}G_p^k, \quad \text{or} \quad [J_0, G_p^{\pm}] = \pm G_p^{\pm}.$$

The product of odd generators are given by the following relation:

$$\{G_{\frac{1}{2}}^j, G_{\frac{1}{2}}^k\} = 2\delta_{j,k}e, \quad \{G_{\frac{-1}{2}}^j, G_{\frac{-1}{2}}^k\} = -2\delta_{j,k}f, \quad \{G_{\frac{-1}{2}}^j, G_{\frac{1}{2}}^k\} = -\delta_{j,k}h - \sqrt{-1}\epsilon_{j,k}J_0,$$

equivalently

$$\begin{aligned} \{G_p^+, G_q^+\} &= \{G_p^-, G_q^-\} = 0, \\ \{G_{\frac{1}{2}}^+, G_{\frac{1}{2}}^-\} &= 2e, \quad \{G_{\frac{-1}{2}}^+, G_{\frac{-1}{2}}^-\} = -2f, \quad \{G_{\frac{-1}{2}}^+, G_{\frac{1}{2}}^-\} = -h - J_0, \quad \{G_{\frac{-1}{2}}^-, G_{\frac{1}{2}}^+\} = -h + J_0. \end{aligned}$$

In conformal field theory, the operators $f, \frac{h}{2}, e, G_{\frac{-1}{2}}^*, G_{\frac{1}{2}}^*, J_0$ are called the momentum, dilation, special-conformal, supersymmetry, s-supersymmetry, and $SO(2)$ -charge generators. Let $\mathfrak{T}^{*1,2}$ be the one dimensional superspace with the coordinate, $t \in \mathfrak{T}^*$, and the two-component Grassmann variable, $\theta = (\theta^1, \theta^2)$. The SCA with $c = 0$ can be represented by the following super-vector fields on $\mathfrak{T}^{*1,2}$,

$$\begin{aligned} L_n &= -t^n(t \frac{\partial}{\partial t} + \frac{n+1}{2}\theta^k \frac{\partial}{\partial \theta^k}), \\ G_p^j &= \sqrt{2\sqrt{-1}}t^{p+\frac{1}{2}}[\theta^j(\frac{\partial}{\partial t} + \frac{2p+1}{2t}\theta^k \frac{\partial}{\partial \theta^k}) - \frac{1}{2\sqrt{-1}}\frac{\partial}{\partial \theta^j}], \\ J_n &= \sqrt{-1}t^n(\theta^1 \frac{\partial}{\partial \theta^2} - \theta^2 \frac{\partial}{\partial \theta^1}). \end{aligned} \tag{25}$$

Under the conformal transformation,

$$(t, \theta) \mapsto (-t^{-1}, -t^{-1}\theta), \quad \theta = (\theta^1, \theta^2),$$

one has the relations,

$$\frac{\partial}{\partial t} \mapsto t^2 \frac{\partial}{\partial t} + t\theta^k \frac{\partial}{\partial \theta^k}, \quad \frac{\partial}{\partial \theta^j} \mapsto -t \frac{\partial}{\partial \theta^j},$$

which give rise to the automorphism (24) of SCA. Let $\mathbb{P}^{1,2}$ be the one-dimensional projective superspace, which is the union of two affine spaces U_0, U_∞ with the super-coordinates:

$$(t, \theta) \in \mathfrak{T}^{1,2} \simeq U_0, \quad (\tilde{t}, \tilde{\theta}) \in \mathfrak{T}^{1,2} \simeq U_\infty.$$

The intersection, $U_0 \cap U_\infty$, consists of elements with $t \neq 0$, or $\tilde{t} \neq 0$, with the coordinate transformation

$$(\tilde{t}, \tilde{\theta}) = (-t^{-1}, -t^{-1}\theta) \in \mathfrak{T}^{*1,2}.$$

We shall denote $\text{Vect}(M)$ the space of super-vector fields of a super-manifold M . Then through (25), one can regard L_n, G_p^j, J_n as elements in $\text{Vect}(U_0 \cap U_\infty)$, and we have

$$\begin{aligned} L_n \quad (n \geq -1), \quad G_p^* \quad (p \geq \frac{-1}{2}), \quad J_m \quad (m \geq 0) &\in \text{Vect}(U_0), \\ L_n \quad (n \leq 1), \quad G_p^* \quad (p \leq \frac{1}{2}), \quad J_m \quad (m \leq 0) &\in \text{Vect}(U_\infty) \end{aligned}$$

Hence

$$\mathfrak{g} = \langle L_n, J_m, G_p^j \rangle_{\mathfrak{T}} \cap \text{Vect}(\mathbb{P}^{1,2}).$$

Now one can clearly see the analogy of above description of N=2 conformal algebra with the usual Virasoro interpretation of vector fields on 1-dimensional projective space $\mathbb{P}^1$.

References

- [1] W. Boucher, D. Friedan and A. Kent, Determinant formulae and unitarity for the $N = 2$ superconformal algebras in two dimensions or exact results on string compactification, Phys. Lett. B 172 (1987) 316-327.
- [2] S.J. Cheng and V. G. Kac, Conformal modules, Asian J. Math. 1 (1997) 181-193.
- [3] V. K. Dobrev, Characters of the unitarizable highest weight modules over $N = 2$ superconformal algebras, Phys. Lett. B 186 (1987) 43-51.
- [4] M. Dözzrapf and B. Gato-Rivera, Determinant formula for the topological $N=2$ superconformal algebra, Nucl. Phys. B558 (1999) 503, hep-th/9905063.
- [5] R. Dijkgraaf, E. Verlinde and H. Verlinde, Topological strings in $d < 1$, Nucl. Phys. B352 (1991) 59.
- [6] T. Eguchi, H. Ooguri, A. Taormina, and S. K. Yang, Superconformal algebras and string compactification on manifolds with $SU(n)$ holonomy, Nucl. Phys. B315 (1989) 193-221.
- [7] M. Eichler and D. Zagier, The theory of Jacobi forms, Progress in Math., Birkhäuser Boston, 1985.
- [8] B.L. Feigin, A.M. Semikhatov, V.A. Sirota, and I. Yu. Tipunin, Resolutions and characters of irreducible representations of the $N = 2$ superconformal algebra, hep-th/9805179.
- [9] D. Gepner, Exactly solvable string compactification on manifolds of $SU(N)$ holonomy, Phys. Lett. 199B (1987) 380; Space-time supersymmetry in compactified string theory and superconformal models, Nucl. Phys. B296 (1988) 757.
- [10] S. J. Gates, Jr. and L. Rana, Superspace Geometrical representations of extended super Virasoro algebra, Phys. Lett. B 438 (1998) 80, hep-th/9806038.
- [11] F. Hirzebruch, T. Berger and R. Jung, Manifolds and modular forms, Aspects of Mathematics, Friedr. Vieweg & Sohn Verlagsgesellschaft mbH, Braunschweig/Wiesbaden, 1992.
- [12] F. Hirzebruch, Complex cobordism and the elliptic genus, (notes by A. Langer), Contemporary Mathematics Vol. 241, 1999 American Mathematical Society, 9-20.
- [13] V. G. Kac, Vertex algebra for beginners, Providence: AMS, University Lecture Notes, vol. 9 (1996).
- [14] T. Kawai, Y. Yamada and S. K. Yang, Elliptic genera and $N = 2$ superconformal field theory, Nucl. Phys. B 414 (1994) 191-212, hep-th/9306096.
- [15] E. B. Kiritsis, Character formulae and the structure of the presentations of the $N = 1, N = 2$ superconformal algebras, Int. J. Mod. Phys. A , 3 (1988) 1871-1906.
- [16] Y. Matsuo, Character formula of $C < 1$ unitary representation of $N = 2$ superconformal algebra, Prog. Theor. Phys. 77 (1987) 793-797.
- [17] F. Ravanini and S-K Yang, Modular invariance in $N = 2$ superconformal field theories, Phys. Lett. B195 (1987), 202-208.

- [18] S. S. Roan, Modular invariance of manifolds with $SU(n)$ holonomy, Intern. J. Math. Vol. 3, No. 3 (1992) 359-395.
- [19] S. S. Roan, Elliptic genus and N=2 conformal algebra (in preparation).
- [20] M. Wakimoto, Fusion rules for N=2 superconformal modules, hep-th/9807144.
- [21] E. Witten, On the Landau-Ginzburg description of $N = 2$ minimal models, Int. J. Mod. Phys. A9 (1994) 4783, hep-th/9304026.